

# ABALOPARATIDE

---

## Products Affected

- TYMLOS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>POSTMENOPAUSAL OSTEOPOROSIS: ONE OF THE FOLLOWING: (1) HIGH RISK FOR FRACTURES DEFINED AS ONE OF THE FOLLOWING: A) HISTORY OF OSTEOPOROTIC (I.E., FRAGILITY, LOW TRAUMA) FRACTURE(S), B) 2 OR MORE RISK FACTORS FOR FRACTURE (E.G., HISTORY OF MULTIPLE RECENT LOW TRAUMA FRACTURES, BMD T-SCORE LESS THAN OR EQUAL TO -2.5, CORTICOSTEROID USE, OR USE OF GNRH ANALOGS), OR C) NO PRIOR TREATMENT FOR OSTEOPOROSIS AND FRAX SCORE OF AT LEAST 20% FOR ANY MAJOR FRACTURE OR OF AT LEAST 3% FOR HIP FRACTURE, (2) UNABLE TO USE ORAL THERAPY (I.E., UPPER GASTROINTESTINAL PROBLEMS UNABLE TO TOLERATE ORAL MEDICATION, LOWER GASTROINTESTINAL PROBLEMS UNABLE TO ABSORB ORAL MEDICATIONS, TROUBLE REMEMBERING TO TAKE ORAL MEDICATIONS OR COORDINATING AN ORAL BISPHOSPHONATE WITH OTHER ORAL MEDICATIONS OR THEIR DAILY ROUTINE), (3) TRIAL OF, INTOLERANCE TO, OR A CONTRAINDICATION TO ONE BISPHOSPHONATE. PATIENT HAS NOT RECEIVED A TOTAL OF 24 MONTHS CUMULATIVE TREATMENT WITH ANY PARATHYROID HORMONE THERAPY.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ABATACEPT IV

## Products Affected

- ORENCIA (WITH MALTOSE)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST.
<b>Coverage Duration</b>	RA, PJIA, PSA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. AGVHD: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. RENEWAL: RA, PJIA, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ABATACEPT SQ

## Products Affected

- ORENCIA
- ORENCIA CLICKJECT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. RENEWAL: RA, PJIA, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ABEMACICLIB

---

## Products Affected

- VERZENIO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ABIRATERONE

---

## Products Affected

- *abiraterone*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC HIGH-RISK CASTRATION-SENSITIVE PROSTATE CANCER, METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: ONE OF THE FOLLOWING: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# ABIRATERONE SUBMICRONIZED

---

## Products Affected

- YONSA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: ONE OF THE FOLLOWING: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ACALABRUTINIB

---

## Products Affected

- CALQUENCE
- CALQUENCE (ACALABRUTINIB MAL)

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ADAGRASIB

---

## Products Affected

- KRAZATI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ADALIMUMAB

## Products Affected

- HUMIRA PEN
- HUMIRA PEN CROHNS-UC-HS START
- HUMIRA PEN PSOR-UVEITS-ADOL HS
- HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML
- HUMIRA(CF)
- HUMIRA(CF) PEDI CROHNS STARTER
- HUMIRA(CF) PEN
- HUMIRA(CF) PEN CROHNS-UC-HS
- HUMIRA(CF) PEN PEDIATRIC UC
- HUMIRA(CF) PEN PSOR-UV-ADOL HS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. UVEITIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OPHTHALMOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PJIA, PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. AS: TRIAL OF OR CONTRAINDICATION TO AN NSAID. PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIEN E, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. CD, UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPURINE, METHOTREXATE, OR MESALAMINE. UVEITIS: NO ISOLATED ANTERIOR UVEITIS. RENEWAL: RA, PJIA, PSA, AS, PSO, HIDRADENITIS SUPPURATIVA, UVEITIS: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# AFATINIB

---

## Products Affected

- GILOTRIF

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS.
<b>Other Criteria</b>	METASTATIC NSCLC WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# AGALSIDASE BETA

## Products Affected

- FABRAZYME

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	FABRY DISEASE: INITIAL: PATIENT IS SYMPTOMATIC OR HAS EVIDENCE OF INJURY FROM GL-3 TO THE KIDNEY, HEART, OR CENTRAL NERVOUS SYSTEM RECOGNIZED BY LABORATORY, HISTOLOGICAL, OR IMAGING FINDINGS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	FABRY DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH NEPHROLOGIST, CARDIOLOGIST, OR SPECIALIST IN GENETICS OR INHERITED METABOLIC DISORDERS.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	FABRY DISEASE: INITIAL: NOT CONCURRENTLY USING AN ALPHA-GAL A PHARMACOLOGICAL CHAPERONE (I.E., GALAFOLD (MIGALASTAT)). RENEWAL: 1) DEMONSTRATED IMPROVEMENT OR STABILIZATION, AND 2) NOT CONCURRENTLY USING AN ALPHA-GAL A PHARMACOLOGICAL CHAPERONE (I.E., GALAFOLD (MIGALASTAT)).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ALECTINIB

---

## Products Affected

- ALECENSA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# ALPELISIB

---

## Products Affected

- PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# AMBRISENTAN

## Products Affected

- *ambrisentan*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: 1) DOCUMENTED CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) OF 3 WOOD UNITS OR GREATER, AND 2) NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PAH: INITIAL: PATIENT DOES NOT HAVE IDIOPATHIC PULMONARY FIBROSIS. RENEWAL: 1) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE, OR 2) A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# AMIVANTAMAB-VMJW

---

## Products Affected

- RYBREVANT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ANAKINRA

---

## Products Affected

- KINERET

PA Criteria	Criteria Details
Exclusion Criteria	CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. ALL OTHERS: 12 MONTHS.
Other Criteria	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, RINVOQ, ENBREL, XELJANZ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# APALUTAMIDE

---

## Products Affected

- ERLEADA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC): TRIAL OF OR CONTRAINDICATION TO XTANDI OR ABIRATERONE. NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): (1) HIGH RISK PROSTATE CANCER (I.E., RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS), AND (2) TRIAL OF OR CONTRAINDICATION TO XTANDI OR NUBEQA. NMCRPC, MCSPC: (1) RECEIVED A BILATERAL ORCHIECTOMY, (2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR (3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE ANALOG. RENEWAL: DIAGNOSIS OF NMCRPC OR MCSPC.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# APOMORPHINE

---

## Products Affected

- *apomorphine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	PD: INITIAL: PHYSICIAN HAS OPTIMIZED DRUG THERAPY FOR PD. RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WHILE ON THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# APOMORPHINE - SL

---

## Products Affected

- KYNMOBI SUBLINGUAL FILM 10 MG, 10-15-20-25-30 MG, 15 MG, 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OF AGE OR OLDER.
<b>Prescriber Restrictions</b>	PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	PD: INITIAL: PHYSICIAN HAS OPTIMIZED DRUG THERAPY FOR PARKINSONS DISEASE. RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WITH THE USE OF KYNMOBI.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# APREMILAST

---

## Products Affected

- OTEZLA
- OTEZLA STARTER

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: MILD PLAQUE PSORIASIS (PSO): ONE OF THE FOLLOWING: 1) PSORIASIS COVERING 2 PERCENT OF BODY SURFACE AREA (BSA), 2) STATIC PHYSICIAN GLOBAL ASSESSMENT (SPGA) SCORE OF 2, OR 3) PSORIASIS AREA AND SEVERITY INDEX (PASI) SCORE OF 2 TO 9. MODERATE TO SEVERE PSO: 1) PSORIASIS COVERING 3 PERCENT OR MORE OF BSA, OR 2) PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. BEHCETS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.



PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. MILD PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL SYSTEMIC AGENT (E.G., METHOTREXATE, ACITRETIN, CYCLOSPORINE) AND ONE CONVENTIONAL TOPICAL AGENT (E.G., PUVA, UVB, TOPICAL CORTICOSTEROIDS). MODERATE TO SEVERE PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. BEHCETS DISEASE: 1) PATIENT HAS ORAL ULCERS OR A HISTORY OF RECURRENT ORAL ULCERS BASED ON CLINICAL SYMPTOMS, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OR MORE CONSERVATIVE TREATMENTS (E.G., COLCHICINE, TOPICAL CORTICOSTEROID, ORAL CORTICOSTEROID). RENEWAL: PSA, PSO, BEHCETS DISEASE: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ASCIMINIB

---

## Products Affected

- SCEMBLIX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML); MUTATIONAL ANALYSIS PRIOR TO INITIATION AND SCEMBLIX IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ASFOTASE

---

## Products Affected

- STRENSIQ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HYPOPHOSPHATASIA (HPP): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST, GENETICIST, OR METABOLIC SPECIALIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: PERINATAL/INFANTILE-ONSET HPP: 1) 6 MONTHS OF AGE OR YOUNGER AT ONSET OF HPP, AND 2) POSITIVE FOR A TISSUE NON-SPECIFIC ALKALINE PHOSPHATASE (TNSALP) (ALPL) GENE MUTATION AS CONFIRMED BY GENETIC TESTING OR TWO OF THE FOLLOWING: (A) SERUM ALKALINE PHOSPHATASE (ALP) LEVEL BELOW THAT OF NORMAL RANGE FOR PATIENT AGE, (B) ELEVATED SERUM PYRIDOXAL-5'-PHOSPHATE (PLP) LEVELS AND NO VITAMIN B6 SUPPLEMENTATION IN THE PREVIOUS WEEK, (C) URINE PHOSPHOETHANOLAMINE (PEA) LEVEL ABOVE THAT OF NORMAL RANGE FOR PATIENT AGE, (D) RADIOGRAPHIC EVIDENCE OF HPP, (E) AT LEAST TWO OF THE FOLLOWING: (I) RACHITIC CHEST DEFORMITY, (II) CRANIOSYNOSTOSIS, (III) DELAY IN SKELETAL GROWTH RESULTING IN DELAY OF MOTOR DEVELOPMENT, (IV) HISTORY OF VITAMIN B6 DEPENDENT SEIZURES, (V) NEPHROCALCINOSIS OR HISTORY OF ELEVATED SERUM CALCIUM, (VI) HISTORY OR PRESENCE OF NON-TRAUMATIC POSTNATAL FRACTURE AND DELAYED FRACTURE HEALING. JUVENILE-ONSET HPP: 1) 18 YEARS OF AGE OR YOUNGER AT ONSET OF HPP, AND 2) POSITIVE FOR A TNSALP ALPL GENE MUTATION AS CONFIRMED BY GENETIC TESTING OR TWO OF THE FOLLOWING: (A) SERUM ALP LEVEL BELOW THAT OF NORMAL RANGE FOR PATIENT AGE, (B) ELEVATED SERUM PLP LEVELS AND NO VITAMIN B6 SUPPLEMENTATION IN THE PREVIOUS WEEK, (C) URINE PEA LEVEL ABOVE THAT OF NORMAL RANGE FOR PATIENT AGE, (D) RADIOGRAPHIC EVIDENCE OF HPP, (E) AT LEAST TWO OF THE FOLLOWING: (I) RACHITIC DEFORMITIES, (II) PREMATURE LOSS OF PRIMARY TEETH PRIOR TO 5 YEARS OF AGE, (III) DELAY IN SKELETAL GROWTH RESULTING IN DELAY OF MOTOR DEVELOPMENT, (IV)</p>

PA Criteria	Criteria Details
	<p>HISTORY OR PRESENCE OF NON-TRAUMATIC FRACTURES OR DELAYED FRACTURE HEALING. ALL INDICATIONS: 1) NOT CURRENTLY RECEIVING TREATMENT WITH A BISPHOSPHONATE, 2) CALCIUM OR PHOSPHATE LEVELS ARE NOT BELOW THE NORMAL RANGE, 3) NOT HAVE A TREATABLE FORM OF RICKETS. RENEWAL: ALL INDICATIONS: 1) IMPROVEMENT IN THE SKELETAL CHARACTERISTICS OF HPP, AND 2) NOT CURRENTLY RECEIVING TREATMENT WITH A BISPHOSPHONATE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ATEZOLIZUMAB

---

## Products Affected

- TECENTRIQ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ATOGEPANT

---

## Products Affected

- QULIPTA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	EPISODIC MIGRAINE PREVENTION: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# AVAPRITINIB

---

## Products Affected

- AYVAKIT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# AVATROMBOPAG

## Products Affected

- DOPTELET (10 TAB PACK)
- DOPTELET (15 TAB PACK)
- DOPTELET (30 TAB PACK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: CHRONIC LIVER DISEASE (CLD): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, GASTROENTEROLOGIST, HEPATOLOGIST, IMMUNOLOGIST, ENDOCRINOLOGIST, OR A SURGEON. CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
Coverage Duration	CLD: 1 MONTH. CHRONIC ITP: INITIAL: 2 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: CLD: 1) PLANNED PROCEDURE 10 TO 13 DAYS AFTER INITIATION OF DOPTELET, AND 2) NOT RECEIVING OTHER THROMBOPOIETIN RECEPTOR AGONISTS (E.G., ROMIPLOSTIM, ELTROMBOPAG, ETC.). CHRONIC ITP: TRIAL OF OR CONTRAINDICATION TO CORTICOSTEROIDS OR IMMUNOGLOBULINS OR INSUFFICIENT RESPONSE TO SPLENECTOMY. RENEWAL: CHRONIC ITP: PATIENT HAD A CLINICAL RESPONSE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# AXITINIB

---

## Products Affected

- INLYTA ORAL TABLET 1 MG, 5 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# AZACITIDINE

---

## Products Affected

- ONUREG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# AZTREONAM

---

## Products Affected

- CAYSTON

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	7 YEARS OF AGE OR OLDER
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BARICITINIB

---

## Products Affected

- OLUMIANT

PA Criteria	Criteria Details
Exclusion Criteria	CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS, SEVERE ALOPECIA AREATA.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS (RA): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	RA: INITIAL: TRIAL OF OR CONTRAINDICATION TO ANY TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BECAPLERMIN

---

## Products Affected

- REGRANEX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	DIABETIC NEUROPATHIC ULCERS: PRESCRIBED BY OR IN CONSULTATION WITH A VASCULAR SURGEON, PODIATRIST, ENDOCRINOLOGIST, PHYSICIAN PRACTICING IN A SPECIALTY WOUND CLINIC OR INFECTIOUS DISEASE SPECIALIST.
<b>Coverage Duration</b>	3 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BEDAQUILINE

---

## Products Affected

- SIRTURO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	24 WEEKS
<b>Other Criteria</b>	PULMONARY MULTI-DRUG RESISTANT TUBERCULOSIS (MDR-TB): SIRTURO USED IN COMBINATION WITH AT LEAST 3 OTHER ANTIBIOTICS FOR THE TREATMENT OF PULMONARY MDR-TB.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BELIMUMAB

## Products Affected

- BENLYSTA INTRAVENOUS
- BENLYSTA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: SYSTEMIC LUPUS ERYTHEMATOSUS (SLE): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. LUPUS NEPHRITIS (LN): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR NEPHROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: SLE: CURRENTLY TAKING CORTICOSTEROIDS, ANTIMALARIALS, NSAIDS, OR IMMUNOSUPPRESSIVE AGENTS. RENEWAL: SLE: PATIENT HAD CLINICAL IMPROVEMENT. LN: IMPROVEMENT IN RENAL RESPONSE FROM BASELINE LABORATORY VALUES (I.E., EGFR OR PROTEINURIA) AND/OR CLINICAL PARAMETERS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# BELUMOSUDIL

---

## Products Affected

- REZUROCK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BELZUTIFAN

---

## Products Affected

- WELIREG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BENDAMUSTINE

---

## Products Affected

- *bendamustine intravenous recon soln*
- BENDEKA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BENRALIZUMAB

---

## Products Affected

- FASENRA
- FASENRA PEN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	ASTHMA: INITIAL: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	ASTHMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.
<b>Coverage Duration</b>	INITIAL: 4 MONTHS, RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>ASTHMA: INITIAL: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE, OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE OF THE FOLLOWING: (A) PATIENT EXPERIENCED AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) PATIENT HAS POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA. RENEWAL: 1) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE, OR (D) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BETAINE

---

## Products Affected

- *betaine*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BEVACIZUMAB-AWWB

---

## Products Affected

- MVASI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BEVACIZUMAB-BVZR

---

## Products Affected

- ZIRABEV

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# BEXAROTENE

---

## Products Affected

- *bexarotene*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BINIMETINIB

---

## Products Affected

- MEKTOVI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BORTEZOMIB

---

## Products Affected

- *bortezomib injection recon soln 1 mg, 2.5 mg* • VELCADE
- BORTEZOMIB INTRAVENOUS RECON SOLN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BOSENTAN

## Products Affected

- TRACLEER ORAL TABLET
- TRACLEER ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: 1) DOCUMENTED CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) OF 3 WOOD UNITS OR GREATER, AND 2) NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PAH: INITIAL: 1) DOES NOT HAVE ELEVATED LIVER ENZYMES (ALT, AST) MORE THAN 3 TIMES UPPER LIMIT OF NORMAL (ULN) OR INCREASE IN BILIRUBIN BY 2 OR MORE TIMES ULN, AND 2) NOT CONCURRENTLY TAKING CYCLOSPORINE A OR GLYBURIDE. RENEWAL: 1) NOT CONCURRENTLY TAKING CYCLOSPORINE A OR GLYBURIDE, 2) AGES 3 TO 17 YEARS OF AGE: A) DEMONSTRATED IMPROVEMENT IN PVR, OR B) REMAINED STABLE OR SHOWN IMPROVEMENT IN EXERCISE ABILITY, 3) AGES 18 YEARS OR OLDER: A) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE, OR B) A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# BOSUTINIB

---

## Products Affected

- BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUSLY TREATED (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND BOSULIF IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# BRIGATINIB

---

## Products Affected

- ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG
- ALUNBRIG ORAL TABLETS,DOSE PACK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# C1 ESTERASE INHIBITOR-CINRYZE

## Products Affected

- CINRYZE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, IMMUNOLOGIST, OR ALLERGIST.
<b>Coverage Duration</b>	INITIAL AND RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT COMPARED TO BASELINE IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# C1 ESTERASE INHIBITOR-HAEGARDA

## Products Affected

- HAEGARDA SUBCUTANEOUS  
RECON SOLN 2,000 UNIT, 3,000 UNIT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, IMMUNOLOGIST, OR ALLERGIST.
<b>Coverage Duration</b>	INITIAL AND RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT COMPARED TO BASELINE IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CABOZANTINIB

---

## Products Affected

- COMETRIQ ORAL CAPSULE 100 MG/DAY(80 MG X1-20 MG X1), 140 MG/DAY(80 MG X1-20 MG X3), 60 MG/DAY (20 MG X 3/DAY)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CABOZANTINIB S-MALATE - CABOMETYX

---

## Products Affected

- CABOMETYX ORAL TABLET 20 MG, 40 MG, 60 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CANAKINUMAB

## Products Affected

- ILARIS (PF)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: CRYOPYRIN-ASSOCIATED PERIODIC SYNDROMES (CAPS), SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA), ADULT-ONSET STILLS DISEASE (AOSD): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR AN IMMUNOLOGIST.
Coverage Duration	INITIAL: AOSD, SJIA: 6 MONTHS, RENEWAL:12 MONTHS. ALL OTHER INDICATIONS: 12 MONTHS.
Other Criteria	INITIAL: AOSD, SJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUGS). RENEWAL: AOSD, SJIA: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CANNABIDIOL

## Products Affected

- EPIDIOLEX

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	DRAVET SYNDROME (DS), LENNOX-GASTAUT SYNDROME (LGS), TUBEROUS SCLEROSIS COMPLEX (TSC): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	INITIAL: 12 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: LENNOX-GASTAUT SYNDROME (LGS): TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM. RENEWAL: DS, LGS, TSC: CONFIRMATION OF DIAGNOSIS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CAPLACIZUMAB YHDP

## Products Affected

- CABLIVI INJECTION KIT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	ACQUIRED THROMBOTIC THROMBOCYTOPENIA PURPURA (ATTP): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST
Coverage Duration	12 MONTHS
Other Criteria	ATTP: CABLIVI WAS PREVIOUSLY INITIATED AS PART OF THE FDA APPROVED TREATMENT REGIMEN IN COMBINATION WITH PLASMA EXCHANGE AND IMMUNOSUPPRESSIVE THERAPY WITHIN AN INPATIENT SETTING. THE PATIENT HAS NOT EXPERIENCED MORE THAN TWO RECURRENCES OF ATTP WHILE ON CABLIVI THERAPY (I.E., NEW DROP IN PLATELET COUNT REQUIRING REPEAT PLASMA EXCHANGE DURING 30 DAYS POST-PLASMA EXCHANGE THERAPY [PEX] AND UP TO 28 DAYS OF EXTENDED THERAPY).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CAPMATINIB

---

## Products Affected

- TABRECTA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CARGLUMIC ACID

## Products Affected

- *carglumic acid*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: ACUTE, CHRONIC HYPERAMMONEMIA (HA) DUE TO N ACETYLGLUTAMATE SYNTHASE (NAGS) DEFICIENCY: NAGS GENE MUTATION IS CONFIRMED BY BIOCHEMICAL OR GENETIC TESTING. ACUTE HA DUE TO PROPIONIC ACIDEMIA (PA): 1) CONFIRMED BY THE PRESENCE OF ELEVATED METHYLCITRIC ACID AND NORMAL METHYLMALONIC ACID, OR 2) GENETIC TESTING CONFIRMING MUTATION IN THE PCCA OR PCCB GENE. ACUTE HA DUE TO METHYLMALONIC ACIDEMIA (MMA): 1) CONFIRMED BY THE PRESENCE OF ELEVATED METHYLMALONIC ACID, METHYLCITRIC ACID, OR 2) GENETIC TESTING CONFIRMING MUTATION IN THE MMUT, MMA, MMAB OR MMADHC GENES.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	ACUTE HA DUE TO NAGS/PA/MMA: 7 DAYS. CHRONIC HA DUE TO NAGS: INITIAL: 6 MOS, RENEWAL: 12 MOS.
<b>Other Criteria</b>	RENEWAL: CHRONIC HA DUE TO NAGS: PATIENT HAS SHOWN CLINICAL IMPROVEMENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# CERITINIB

---

## Products Affected

- ZYKADIA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CERTOLIZUMAB PEGOL

## Products Affected

- CIMZIA
- CIMZIA POWDER FOR RECONST

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, TREMFYA, XELJANZ, RINVOQ, SKYRIZI. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, SKYRIZI. NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, RINVOQ. PATIENTS WHO ARE PREGNANT, BREASTFEEDING, OR TRYING TO BECOME PREGNANT ARE EXCLUDED FROM STEP CRITERIA FOR ALL INDICATIONS. RENEWAL: RA, PSA, AS, PSO, NR-AXSPA: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CETUXIMAB

---

## Products Affected

- ERBITUX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CLADRIBINE

## Products Affected

- MAVENCLAD (10 TABLET PACK)
- MAVENCLAD (4 TABLET PACK)
- MAVENCLAD (5 TABLET PACK)
- MAVENCLAD (6 TABLET PACK)
- MAVENCLAD (7 TABLET PACK)
- MAVENCLAD (8 TABLET PACK)
- MAVENCLAD (9 TABLET PACK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 48 WEEKS.
Other Criteria	MS: INITIAL: HAS NOT RECEIVED A TOTAL OF TWO YEARS OF MAVENCLAD TREATMENT (I.E., TWO YEARLY TREATMENT COURSES OF 2 CYCLES IN EACH). RENEWAL: 1) HAS DEMONSTRATED CLINICAL BENEFIT COMPARED TO PRE-TREATMENT BASELINE, 2) DOES NOT HAVE LYMPHOPENIA, AND 3) HAS NOT RECEIVED A TOTAL OF TWO YEARS OF MAVENCLAD TREATMENT (I.E., TWO YEARLY TREATMENT COURSES OF 2 CYCLES IN EACH).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# CLOBAZAM-SYMPAZAN

---

## Products Affected

- SYMPAZAN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	LENNOX-GASTAUT SYNDROME (LGS): THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	LGS: 1) PATIENT IS UNABLE TO TAKE TABLETS OR SUSPENSION, AND 2) TRIAL OF OR CONTRAINDICATION TO A FORMULARY CLOBAZAM AGENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# COBIMETINIB

---

## Products Affected

- COTELLIC

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# COLCHICINE

---

## Products Affected

- *colchicine oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	PROPHYLAXIS OF GOUT FLARES: 16 YEARS AND OLDER
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PROPHYLAXIS OF GOUT FLARES: TRIAL OF OR CONTRAINDICATION TO COLCHICINE CAPSULES (MITIGARE) IF AGE 18 YEARS OR OLDER.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# CORTICOTROPIN

## Products Affected

- ACTHAR
- CORTROPHIN GEL

PA Criteria	Criteria Details
Exclusion Criteria	INITIAL: NOT APPROVED FOR DIAGNOSTIC PURPOSES.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MULTIPLE SCLEROSIS (MS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, ALLERGIST/IMMUNOLOGIST, OPHTHALMOLOGIST, PULMONOLOGIST OR NEPHROLOGIST.
Coverage Duration	INFANTILE SPASMS AND MS: 28 DAYS. OTHER FDA APPROVED INDICATIONS: INITIAL AND RENEWAL: 28 DAYS
Other Criteria	INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS: TRIAL OF OR CONTRAINDICATION TO INTRAVENOUS (IV) CORTICOSTEROIDS. ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MS: TRIAL OF OR CONTRAINDICATION TO A STANDARD OF CARE THERAPY. RENEWAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MS: 1) DEMONSTRATED CLINICAL BENEFIT WHILE ON THERAPY AS INDICATED BY SYMPTOM RESOLUTION AND/OR NORMALIZATION OF LABORATORY TESTS, AND 2) CONTINUES TO POSSESS CONTRAINDICATION TO IV CORTICOSTEROIDS. PART B BEFORE PART D STEP THERAPY, APPLIES ONLY TO BENEFICIARIES IN AN MA-PD PLAN.
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CRIZANLIZUMAB-TMCA

## Products Affected

- ADAKVEO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	SICKLE CELL DISEASE (SCD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST
<b>Coverage Duration</b>	INITIAL: 12 MONTHS. RENEWAL: LIFETIME
<b>Other Criteria</b>	SCD: INITIAL: ADULTS (18 YEARS OR OLDER): ONE OF THE FOLLOWING: (1) AT LEAST 2 SICKLE CELL CRISES IN THE PAST YEAR, (2) SICKLE-CELL ASSOCIATED SYMPTOMS WHICH ARE INTERFERING WITH ACTIVITIES OF DAILY LIVING, OR (3) HISTORY OF OR HAS RECURRENT ACUTE CHEST SYNDROME (ACS). PATIENTS BETWEEN THE AGES OF 16 TO 17 YEARS, WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA. RENEWAL FOR ALL PATIENTS: MAINTAINED OR EXPERIENCED A REDUCTION IN ACUTE COMPLICATIONS OF SICKLE CELL DISEASE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CRIZOTINIB

---

## Products Affected

- XALKORI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# CYSTEAMINE HYDROCHLORIDE

---

## Products Affected

- CYSTADROPS
- CYSTARAN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DABRAFENIB

---

## Products Affected

- TAFINLAR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DACOMITINIB

---

## Products Affected

- VIZIMPRO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	METASTATIC NSCLC: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DALFAMPRIDINE

---

## Products Affected

- *dalfampridine*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	MULTIPLE SCLEROSIS (MS): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	MS: INITIAL: HAS SYMPTOMS OF A WALKING DISABILITY SUCH AS MILD TO MODERATE BILATERAL LOWER EXTREMITY WEAKNESS OR UNILATERAL WEAKNESS PLUS LOWER EXTREMITY OR TRUNCAL ATAXIA. RENEWAL: IMPROVEMENT IN WALKING ABILITY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# DARATUMUMAB

---

## Products Affected

- DARZALEX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DARATUMUMAB-HYALURONIDASE-FIHJ

---

## Products Affected

- DARZALEX FASPRO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DAROLUTAMIDE

## Products Affected

- NUBEQA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL AND RENEWAL: 12 MONTHS.
Other Criteria	<p>INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): PATIENT HAS HIGH RISK PROSTATE CANCER (I.E., RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS) AND ONE OF THE FOLLOWING: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. METASTATIC HORMONE-SENSITIVE PROSTATE CANCER (MHSPC): 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: NMCRPC: NO ADDITIONAL CRITERIA REQUIRED. MHSPC: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.</p>

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DASATINIB

---

## Products Affected

- SPRYCEL ORAL TABLET 100 MG, 140 MG, 20 MG, 50 MG, 70 MG, 80 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	PREVIOUSLY TREATED Ph+ CML: MUTATIONAL ANALYSIS PRIOR TO INITIATION AND SPRYCEL IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DECITABINE/CEDAZURIDINE

---

## Products Affected

- INQOVI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DEFERASIROX

## Products Affected

- *deferasirox*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 1000 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). NON-TRANSFUSION DEPENDENT THALASSEMIA (NTDT): 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS) AND 2) LIVER IRON CONCENTRATION (LIC) OF 5 MG FE/G DRY WEIGHT OR GREATER. RENEWAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 500 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). NTDT: 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS) OR 2) LIC OF 3 MG FE/G DRY WEIGHT OR GREATER.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: CHRONIC IRON OVERLOAD: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL FOR ALL INDICATIONS: FORMULARY VERSION OF DEFERASIROX SPRINKLE: TRIAL OF OR CONTRAINDICATION TO GENERIC DEFERASIROX TABLET OR TABLET FOR ORAL SUSPENSION.
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# DEFERIPRONE

## Products Affected

- *deferiprone*
- FERRIPROX (2 TIMES A DAY)
- FERRIPROX ORAL SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	TRANSFUSIONAL IRON OVERLOAD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: TRANSFUSIONAL IRON OVERLOAD DUE TO THALASSEMIA SYNDROMES: 1) TRIAL OF, CONTRAINDICATION, INTOLERABLE TOXICITIES, OR CLINICALLY SIGNIFICANT ADVERSE EFFECTS TO ONE OF THE FOLLOWING: FORMULARY VERSION OF DEFERASIROX OR DEFEROXAMINE, OR 2) CURRENT CHELATION THERAPY (I.E., FORMULARY VERSION OF DEFERASIROX OR DEFEROXAMINE) IS INADEQUATE. TRANSFUSIONAL IRON OVERLOAD DUE TO SICKLE CELL DISEASE OR OTHER ANEMIAS: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF DEFERASIROX OR DEFEROXAMINE. RENEWAL (ALL INDICATIONS): SERUM FERRITIN LEVELS CONSISTENTLY ABOVE 500MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS).
Indications	All FDA-approved Indications.
Off Label Uses	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# DEFEROXAMINE

## Products Affected

- *deferoxamine*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	CHRONIC IRON OVERLOAD: 3 YEARS OR OLDER
<b>Prescriber Restrictions</b>	CHRONIC IRON OVERLOAD: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	CHRONIC IRON OVERLOAD: INITIAL: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 1000MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). RENEWAL: SERUM FERRITIN LEVELS MUST BE CONSISTENTLY ABOVE 500MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# DENOSUMAB-XGEVA

---

## Products Affected

- XGEVA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DEUTETRABENAZINE

---

## Products Affected

- AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HUNTINGTON DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST. TARDIVE DYSKINESIA: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR MOVEMENT DISORDER SPECIALIST.
Coverage Duration	12 MONTHS
Other Criteria	TARDIVE DYSKINESIA: PATIENT HAS A HISTORY OF USING AGENTS THAT CAUSE TARDIVE DYSKINESIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DICLOFENAC GEL

---

## Products Affected

- *diclofenac sodium topical gel 3%*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DICLOFENAC TOPICAL SOLUTION

---

## Products Affected

- *diclofenac sodium topical solution in metered-dose pump*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	OSTEOARTHRITIS OF THE KNEE: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF DICLOFENAC SODIUM 1% TOPICAL GEL AND A FORMULARY VERSION OF DICLOFENAC SODIUM 1.5% TOPICAL DROPS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# DIMETHYL FUMARATE

---

## Products Affected

- *dimethyl fumarate oral capsule, delayed release(dr/lec) 120 mg, 120 mg (14)- 240 mg (46), 240 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DIROXIMEL FUMARATE

---

## Products Affected

- VUMERITY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DOSTARLIMAB-GXLY

---

## Products Affected

- JEMPERLI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DRONABINOL

---

## Products Affected

- *dronabinol*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY: TRIAL OF OR CONTRAINDICATION TO ONE ANTIEMETIC THERAPY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D FOR THE INDICATION OF NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# DROXIDOPA

## Products Affected

- *droxidopa*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	NEUROGENIC ORTHOSTATIC HYPOTENSION (NOH): INITIAL: 1) BASELINE BLOOD PRESSURE READINGS WHILE THE PATIENT IS SITTING AND ALSO WITHIN 3 MINUTES OF STANDING FROM A SUPINE POSITION. 2) A DECREASE OF AT LEAST 20 MMHG IN SYSTOLIC BLOOD PRESSURE OR 10 MMHG DIASTOLIC BLOOD PRESSURE WITHIN THREE MINUTES AFTER STANDING FROM A SITTING POSITION.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NOH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR CARDIOLOGIST.
<b>Coverage Duration</b>	INITIAL: 3 MONTHS RENEWAL: 12 MONTHS
<b>Other Criteria</b>	NOH: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# DUPILUMAB

## Products Affected

- DUPIXENT PEN
- DUPIXENT SYRINGE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: EOSINOPHILIC ASTHMA: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS. EOSINOPHILIC ESOPHAGITIS (EOE): DIAGNOSIS CONFIRMED BY ESOPHAGOGASTRODUODENOSCOPY (EGD) WITH BIOPSY.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: AD, PN: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE. CRSWNP: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. EOE: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, ALLERGIST, OR IMMUNOLOGIST.
<b>Coverage Duration</b>	INITIAL: AD, CRSWNP, EOE, PN: 6 MOS, ASTHMA: 4 MOS. RENEWAL: ALL INDICATIONS: 12 MOS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: AD: 1) AD COVERING AT LEAST 10 PERCENT OF BODY SURFACE AREA OR AD AFFECTING THE FACE, HEAD, NECK, HANDS, FEET, GROIN, OR INTERTRIGINOUS AREAS, 2) INTRACTABLE PRURITUS OR CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, 3) TRIAL OF OR CONTRAINDICATION TO ONE TOPICAL (CORTICOSTEROID, CALCINEURIN INHIBITOR, PDE4 INHIBITOR, OR JAK INHIBITOR), AND 4) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGIC/JAK INHIBITOR FOR AD. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NO CONCURRENT USE WITH XOLAIR OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA. CRSWNP: 1) EVIDENCE OF NASAL POLYPS BY DIRECT EXAMINATION, ENDOSCOPY OR SINUS CT SCAN, 2) INADEQUATELY CONTROLLED DISEASE AS DETERMINED BY USE OF SYSTEMIC STEROIDS IN THE PAST 2 YEARS OR ENDOSCOPIC SINUS SURGERY, AND 3) A 90 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID. PN: 1) CHRONIC PRURITIS (ITCH MORE</p>

PA Criteria	Criteria Details
	<p>THAN 6 WEEKS), MULTIPLE PRURIGINOUS LESIONS, AND HISTORY OR SIGN OF A PROLONGED SCRATCHING BEHAVIOR, 2) TRIAL OF OR CONTRAINDICATION TO ONE TOPICAL (CORTICOSTEROID OR CALCIPOTRIOL). RENEWAL: AD: 1) IMPROVEMENT WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGIC/JAK INHIBITOR FOR AD. CRSWNP, EOE: IMPROVEMENT WHILE ON THERAPY. ASTHMA: 1) NO CONCURRENT USE WITH XOLAIR, OR OTHER ANTI-IL5 BIOLOGICS FOR ASTHMA, 2) CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE, OR (D) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS. PN: IMPROVEMENT OR REDUCTION OF PRURITIS OR PRURIGINOUS LESIONS.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# DUVELISIB

---

## Products Affected

- COPIKTRA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# EDARAVONE

## Products Affected

- RADICAVA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	AMYOTROPHIC LATERAL SCLEROSIS (ALS): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR ALS SPECIALIST AT AN ALS SPECIALTY CENTER OR CARE CLINIC.
<b>Coverage Duration</b>	ALS: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	ALS: INITIAL: 1) DURATION OF DISEASE (FROM ONSET OF SYMPTOMS) IS LESS THAN OR EQUAL TO 2 YEARS, 2) NORMAL RESPIRATORY FUNCTION, 3) HAS MILD TO MODERATE ALS WITH A SCORE OF 2 OR HIGHER IN ALL OF THE FOLLOWING 12 ITEMS OF THE AMYOTROPHIC LATERAL SCLEROSIS FUNCTIONAL RATING SCALE REVISED (ALSFRS-R): SPEECH, SALIVATION, SWALLOWING, HANDWRITING, CUTTING FOOD, DRESSING AND HYGIENE, TURNING IN BED, WALKING, CLIMBING STAIRS, DYSPNEA, ORTHOPNEA, RESPIRATORY INSUFFICIENCY, AND 4) TRIAL OF RILUZOLE TABLET OR CURRENTLY TAKING RILUZOLE TABLET. RENEWAL: 1) DOES NOT REQUIRE INVASIVE VENTILATION, AND 2) HAS IMPROVED BASELINE FUNCTIONAL ABILITY OR HAS MAINTAINED A SCORE OF 2 OR HIGHER IN ALL 12 ITEMS OF THE ALSFRS-R.
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# EFLAPEGRASTIM-XNST

---

## Products Affected

- ROLVEDON

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: NYVEPRIA. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ELAGOLIX SODIUM

## Products Affected

- ORILISSA ORAL TABLET 150 MG, 200 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
<b>Age Restrictions</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 18 YEARS OF AGE OR OLDER.
<b>Prescriber Restrictions</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
<b>Coverage Duration</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 2) TRIAL OF OR CONTRAINDICATION TO A NSAID AND PROGESTIN-CONTAINING PREPARATION. RENEWAL: 1) IMPROVEMENT IN PAIN ASSOCIATED WITH ENDOMETRIOSIS WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# ELAPEGADEMASE-LVLR

## Products Affected

- REVCOVI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	ADENOSINE DEAMINASE SEVERE COMBINED IMMUNE DEFICIENCY (ADA-SCID): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH IMMUNOLOGIST, HEMATOLOGIST/ONCOLOGIST, OR PHYSICIAN SPECIALIZING IN INHERITED METABOLIC DISORDERS
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	ADA-SCID: INITIAL: ADA-SCID AS MANIFESTED BY ONE OF THE FOLLOWING: (1) CONFIRMATORY GENETIC TEST OR (2) SUGGESTIVE LABORATORY FINDINGS (E.G., ELEVATED DEOXYADENOSINE NUCLEOTIDE [DAXP] LEVELS, LYMPHOPENIA) AND HALLMARK SIGNS/SYMPTOMS (E.G., RECURRENT INFECTIONS, FAILURE TO THRIVE, PERSISTENT DIARRHEA). RENEWAL: 1) IMPROVEMENT OR MAINTENANCE OF IMMUNE FUNCTION FROM BASELINE, AND 2) HAS NOT RECEIVED SUCCESSFUL HCT OR GENE THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ELEXACAFITOR-TEZACAFITOR-IVACAFITOR

## Products Affected

- TRIKAFTA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	CYSTIC FIBROSIS (CF); INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: LIFETIME.
<b>Other Criteria</b>	CF: RENEWAL: 1) MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR 2) REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# ELIGLUSTAT

---

## Products Affected

- CERDELGA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ELOSULFASE ALFA

---

## Products Affected

- VIMIZIM

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ELTROMBOPAG

## Products Affected

- PROMACTA ORAL POWDER IN PACKET 12.5 MG, 25 MG
- PROMACTA ORAL TABLET 12.5 MG, 25 MG, 50 MG, 75 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: ITP: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
Coverage Duration	ITP: INITIAL: 2 MO, RENEWAL: 12 MO. HEPATITIS C, SEVERE APLASTIC ANEMIA: 12 MO.
Other Criteria	INITIAL: PERSISTENT OR CHRONIC IMMUNE THROMBOCYTOPENIA PURPURA (ITP): TRIAL OF OR CONTRAINDICATION TO CORTICOSTEROIDS, IMMUNOGLOBULINS, OR AN INSUFFICIENT RESPONSE TO SPLENECTOMY. ALL INDICATIONS: APPROVAL FOR PROMACTA ORAL SUSPENSION PACKETS REQUIRES A TRIAL OF PROMACTA TABLETS OR PATIENT IS UNABLE TO TAKE TABLET FORMULATION. RENEWAL: ITP: PATIENT HAS SHOWN A CLINICAL RESPONSE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# EMAPALUMAB-LZSG

## Products Affected

- GAMIFANT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS (HLH): INITIAL: 1) A GENETIC TEST IDENTIFYING HLH-ASSOCIATED GENE MUTATION (E.G., PRF1, UNC13D), OR 2) HAS AT LEAST FIVE OF THE FOLLOWING EIGHT DIAGNOSTIC CRITERIA FOR HLH: (A) FEVER, (B) SPLENOMEGALY, (C) CYTOPENIAS (AFFECTING AT LEAST 2 OF 3 CELL LINEAGES), (D) HYPERTRIGLYCERIDEMIA OR HYPOFIBRINOGENEMIA, (E) HEMOPHAGOCYTOSIS IN BONE MARROW OR SPLEEN OR LYMPH NODES AND NO EVIDENCE OF MALIGNANCY, (F) LOW OR ABSENT NATURAL KILLER-CELL ACTIVITY, (G) FERRITIN LEVEL OF 500 MCG/L OR GREATER, (H) SOLUBLE CD25 LEVEL OF 2,400 U/ML OR GREATER.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HLH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN IMMUNOLOGIST, HEMATOLOGIST, OR ONCOLOGIST.
<b>Coverage Duration</b>	INITIAL AND RENEWAL: 8 WEEKS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	HLH: INITIAL: 1) CONCURRENT THERAPY WITH DEXAMETHASONE, AND 2) ONE OF THE FOLLOWING: (A) HAS REFRACTORY, RECURRENT, OR PROGRESSIVE DISEASE, OR (B) HAD A TRIAL OF OR INTOLERANCE TO CONVENTIONAL HLH THERAPY (I.E., CHEMOTHERAPY, STEROIDS, IMMUNOTHERAPY). RENEWAL: 1) HAS NOT RECEIVED SUCCESSFUL HEMATOPOIETIC STEM CELL TRANSPLANTATION, AND 2) DEMONSTRATED IMPROVED IMMUNE SYSTEM RESPONSE FROM BASELINE (E.G., RESOLUTION OF FEVER, DECREASED SPLENOMEGALY, IMPROVEMENT IN CNS SYMPTOMS, IMPROVED CBC, INCREASED FIBRINOGEN LEVELS, REDUCED D-DIMER, REDUCED FERRITIN, REDUCED SOLUBLE CD25 LEVELS.)
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ENASIDENIB

---

## Products Affected

- IDHIFA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ENCORAFENIB

---

## Products Affected

- BRAFTOVI ORAL CAPSULE 75 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ENTRECTINIB

---

## Products Affected

- ROZLYTREK ORAL CAPSULE 100 MG, 200 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# ENZALUTAMIDE

## Products Affected

- XTANDI ORAL CAPSULE
- XTANDI ORAL TABLET 40 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	INITIAL: CASTRATION-RESISTANT PROSTATE CANCER (CRPC) THAT IS NOT METASTATIC: PATIENT HAS HIGH RISK PROSTATE CANCER (I.E. RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). CRPC (INCLUDES NON-METASTATIC AND METASTATIC) OR METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC): 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, OR 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: DIAGNOSIS OF CRPC (INCLUDES NON-METASTATIC AND METASTATIC) OR MCSPC.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# EPOETIN ALFA-EPBX

## Products Affected

- RETACRIT INJECTION SOLUTION  
10,000 UNIT/ML, 2,000 UNIT/ML,  
20,000 UNIT/2 ML, 20,000 UNIT/ML,  
3,000 UNIT/ML, 4,000 UNIT/ML, 40,000  
UNIT/ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: CHRONIC KIDNEY DISEASE (CKD), ANEMIA RELATED TO ZIDOVUDINE, OR CANCER CHEMOTHERAPY: HEMOGLOBIN LEVEL OF LESS THAN 10G/DL. ELECTIVE, NON-CARDIAC, NON-VASCULAR SURGERY: HEMOGLOBIN LEVEL LESS THAN 13G/DL. RENEWAL: CKD: 1) HEMOGLOBIN LEVEL IS LESS THAN 10G/DL, OR 2) HEMOGLOBIN LEVEL HAS REACHED 10G/DL AND DOSE REDUCTION/INTERRUPTION IS REQUIRED TO REDUCE THE NEED FOR BLOOD TRANSFUSIONS. ANEMIA RELATED TO ZIDOVUDINE: HEMOGLOBIN LEVEL BETWEEN 10G/DL AND 12G/DL. CANCER CHEMOTHERAPY: 1) HEMOGLOBIN LEVEL OF LESS THAN 10 G/DL, OR 2) THE HEMOGLOBIN LEVEL DOES NOT EXCEED A LEVEL NEEDED TO AVOID RBC TRANSFUSION.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	ANEMIA FROM CHEMO/CKD WITHOUT DIALYSIS/ZIDOVUDINE: INITIAL/RENEWAL: 12 MONTHS. SURGERY: 1 MONTH.
<b>Other Criteria</b>	RENEWAL: CKD: NOT RECEIVING DIALYSIS TREATMENT. THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# EPOPROSTENOL IV

## Products Affected

- *epoprostenol (glycine)*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: (1) DOCUMENTED CONFIRMATORY PAH DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER, AND (2) NYHA-WHO FUNCTIONAL CLASS III-IV SYMPTOMS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL AND RENEWAL: 12 MONTHS
<b>Other Criteria</b>	PAH: RENEWAL: IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ERDAFITINIB

---

## Products Affected

- BALVERSA ORAL TABLET 3 MG, 4 MG, 5 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ERLOTINIB

---

## Products Affected

- *erlotinib oral tablet 100 mg, 150 mg, 25 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	METASTATIC NSCLC WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ESKETAMINE

## Products Affected

- SPRAVATO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: TREATMENT-RESISTANT DEPRESSION (TRD), MAJOR DEPRESSIVE DISORDER (MDD): PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST.
Coverage Duration	INITIAL: TRD: 3 MONTHS. MDD: 4 WEEKS. RENEWAL: TRD, MDD: 12 MONTHS.
Other Criteria	INITIAL: TRD: 1) NON-PSYCHOTIC, UNIPOLAR DEPRESSION, 2) NO ACTIVE SUBSTANCE ABUSE, AND 3) ADEQUATE TRIAL (AT LEAST 4 WEEKS) OF AT LEAST TWO ANTIDEPRESSANT AGENTS FROM DIFFERENT CLASSES THAT ARE INDICATED FOR DEPRESSION. MDD: 1) NON-PSYCHOTIC, UNIPOLAR DEPRESSION, AND 2) NO ACTIVE SUBSTANCE ABUSE. RENEWAL: TRD, MDD: DEMONSTRATED CLINICAL BENEFIT (IMPROVEMENT IN DEPRESSION) COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ETANERCEPT

---

## Products Affected

- ENBREL
- ENBREL MINI
- ENBREL SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE OR GENITAL AREA.
Age Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), PSORIATIC ARTHRITIS (PSA): 18 YEARS OR OLDER.
Prescriber Restrictions	INITIAL: RA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), AS: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSA: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.



PA Criteria	Criteria Details
<b>Other Criteria</b>	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE: IS REQUIRED. PJIA, PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. AS: TRIAL OF OR CONTRAINDICATION TO AN NSAID. PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. RENEWAL: RA, PJIA, PSA, AS, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ETEPLIRSEN

## Products Affected

- EXONDYS-51

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	DUCHENNE MUSCULAR DYSTROPHY (DMD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	DMD: INITIAL: PATIENT IS AMBULATORY AND CURRENTLY RECEIVING TREATMENT WITH OR HAS A CONTRAINDICATION TO CORTICOSTEROIDS. RENEWAL: MAINTAINED OR DEMONSTRATED A LESS THAN EXPECTED DECLINE IN AMBULATORY ABILITY IN MUSCLE FUNCTION ASSESSMENTS OR OTHER MUSCLE FUNCTION (I.E., PULMONARY OR CARDIAC FUNCTION). THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# EVEROLIMUS

---

## Products Affected

- *everolimus (antineoplastic) oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg*
- *everolimus (antineoplastic) oral tablet for suspension*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADVANCED RENAL CELL CARCINOMA (RCC): TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF SUNITINIB OR SORAFENIB.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FEDRATINIB

---

## Products Affected

- INREBIC

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MYELOFIBROSIS: INITIAL: TRIAL OF OR CONTRAINDICATION TO JAKAFI (RUXOLITINIB). RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# FENFLURAMINE

## Products Affected

- FINTEPLA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: DRAVET SYNDROME, LENNOX-GASTAUT SYNDROME (LGS): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	DRAVET SYNDROME: INITIAL/RENEWAL: 12 MONTHS. LGS: 12 MONTHS.
Other Criteria	INITIAL: LGS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM. RENEWAL: DRAVET SYNDROME: PATIENT HAS SHOWN CONTINUED CLINICAL BENEFIT (E.G. REDUCTION OF SEIZURES, REDUCED LENGTH OF SEIZURES, SEIZURE CONTROL MAINTAINED).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FENTANYL CITRATE

---

## Products Affected

- *fentanyl citrate buccal lozenge on a handle*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	CANCER RELATED PAIN: 1) CURRENTLY ON A MAINTENANCE DOSE OF CONTROLLED-RELEASE OPIOID PAIN MEDICATION, AND 2) TRIAL OF OR CONTRAINDICATION TO AT LEAST ONE IMMEDIATE-RELEASE ORAL OPIOID PAIN AGENT OR PATIENT HAS DIFFICULTY SWALLOWING TABLETS/CAPSULES. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# FILGRASTIM-AAFI

---

## Products Affected

- NIVESTYM

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# FILGRASTIM-AYOW

---

## Products Affected

- RELEUKO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT NIVESTYM, WHERE INDICATIONS ALIGN.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# FILGRASTIM-SNDZ

---

## Products Affected

- ZARXIO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: NIVESTYM.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# FINASTERIDE/TADALAFIL

---

## Products Affected

- ENTADFI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	BENIGN PROSTATIC HYPERPLASIA (BPH): 18 YEARS OF AGE OR OLDER
Prescriber Restrictions	
Coverage Duration	26 WEEKS
Other Criteria	BPH: 1) TRIAL OF OR CONTRAINDICATION TO ONE 5-ALPHA-REDUCTASE INHIBITOR, AND 2) TRIAL OF OR CONTRAINDICATION TO TADALAFIL 2.5MG OR TADALAFIL 5MG.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FINERENONE

---

## Products Affected

- KERENDIA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# FINGOLIMOD

---

## Products Affected

- *fingolimod*
- GILENYA ORAL CAPSULE 0.25 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FINGOLIMOD LAURYL SULFATE

---

## Products Affected

- TASCENSO ODT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	MULTIPLE SCLEROSIS (MS): (1) UNABLE TO SWALLOW FINGOLIMOD CAPSULES, AND (2) TRIAL OF OR CONTRAINDICATION TO FINGOLIMOD CAPSULES.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# FOSTAMATINIB

---

## Products Affected

- TAVALISSE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
Coverage Duration	INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	ITP: RENEWAL: PATIENT HAS SHOWN A CLINICAL RESPONSE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FREMANEZUMAB-VFRM

## Products Affected

- AJOVY AUTOINJECTOR
- AJOVY SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	MIGRAINE PREVENTION: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# FUTIBATINIB

---

## Products Affected

- LYTGOBI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INTRAHEPATIC CHOLANGIOCARCINOMA (ICCA): COMPLETE A COMPREHENSIVE OPHTHALMOLOGICAL EXAMINATION, INCLUDING OPTICAL COHERENCE TOMOGRAPHY (OCT), PRIOR TO THE INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# GALCANEZUMAB-GNLM

## Products Affected

- EMGALITY PEN 3)
- EMGALITY SYRINGE  
SUBCUTANEOUS SYRINGE 120  
MG/ML, 300 MG/3 ML (100 MG/ML X

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: MIGRAINE PREVENTION: 6 MOS. EPISODIC CLUSTER HEADACHE: 3 MOS. RENEWAL (ALL): 12 MOS.
Other Criteria	INITIAL: MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL. RENEWAL: MIGRAINE PREVENTION: 1) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY, AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. EPISODIC CLUSTER HEADACHE: IMPROVEMENT IN EPISODIC CLUSTER HEADACHE FREQUENCY AS COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# GANAXOLONE

---

## Products Affected

- ZTALMY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# GEFITINIB

---

## Products Affected

- IRESSA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	METASTATIC NSCLC WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# GILTERITINIB

---

## Products Affected

- XOSPATA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# GIVOSIRAN

## Products Affected

- GIVLAARI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	ACUTE HEPATIC PORPHYRIA (AHP): INITIAL: GENETIC CONFIRMATION OF MUTATION OR ELEVATED URINARY OR PLASMA PBG (PORPHOBILINOGEN) OR ALA (AMINOLEVULINIC ACID).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	ACUTE HEPATIC PORPHYRIA (AHP): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A GENETICIST, HEPATOLOGIST, HEMATOLOGIST, GASTROENTEROLOGIST, NEUROLOGIST, DERMATOLOGIST, OR A HEALTHCARE PROVIDER EXPERIENCED IN MANAGING AHP.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS
<b>Other Criteria</b>	AHP: INITIAL: EXPERIENCED TWO OR MORE AHP ATTACKS IN THE PAST 12 MONTHS. RENEWAL: 1) ACHIEVED OR MAINTAINED CLINICAL BENEFIT COMPARED TO BASELINE, AND 2) HAS NOT RECEIVED A LIVER TRANSPLANT. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# GLASDEGIB

---

## Products Affected

- DAURISMO ORAL TABLET 100 MG,  
25 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# GLATIRAMER

---

## Products Affected

- COPAXONE SUBCUTANEOUS SYRINGE 20 MG/ML, 40 MG/ML
- *glatiramer subcutaneous syringe 20 mg/ml, 40 mg/ml*
- *glatopa subcutaneous syringe 20 mg/ml, 40 mg/ml*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# GLECAPREVIR/PIBRENTASVIR

## Products Affected

- MAVYRET ORAL TABLET

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HCV RNA LEVEL WITHIN PAST 6 MONTHS
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
<b>Other Criteria</b>	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) TRIAL OF A PREFERRED FORMULARY ALTERNATIVE INCLUDING HARVONI OR EPCLUSA WHEN THESE AGENTS ARE CONSIDERED ACCEPTABLE FOR TREATMENT OF THE SPECIFIC GENOTYPE PER AASLD/IDSA GUIDANCE, 3) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: CARBAMAZEPINE, RIFAMPIN, ETHINYL ESTRADIOL-CONTAINING MEDICATION, ATAZANAVIR, DARUNAVIR, LOPINAVIR, RITONAVIR, EFAVIRENZ, ATORVASTATIN, LOVASTATIN, SIMVASTATIN, ROSUVASTATIN AT DOSES GREATER THAN 10MG, CYCLOSPORINE AT DOSES GREATER THAN 100MG PER DAY, EPCLUSA, HARVONI, VOSEVI, OR ZEPATIER, 4) PATIENT MUST NOT HAVE PRIOR FAILURE OF A DAA REGIMEN WITH NS5A INHIBITOR AND HCV PROTEASE INHIBITOR, AND 5) DOES NOT HAVE MODERATE OR SEVERE HEPATIC IMPAIRMENT (CHILD PUGH B OR C).
<b>Indications</b>	All FDA-approved Indications.



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# GLYCEROL PHENYL BUTYRATE

---

## Products Affected

- RAVICTI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	UREA CYCLE DISORDER (UCD): INITIAL: DIAGNOSIS IS CONFIRMED BY ENZYMATIC, BIOCHEMICAL OR GENETIC TESTING
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS
Other Criteria	UCD: INITIAL: TRIAL OF OR CONTRAINDICATION TO SODIUM PHENYL BUTYRATE. RENEWAL: PATIENT HAS CLINICAL BENEFIT FROM BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# GOSERELIN

## Products Affected

- ZOLADEX

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
<b>Coverage Duration</b>	STAGE B2-C PROSTATIC CARCINOMA: 4 MOS. ENDOMETRIOSIS: 6 MOS PER LIFETIME. ALL OTHERS: 12 MONTHS.
<b>Other Criteria</b>	ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 6 MONTHS OF TREATMENT PER LIFETIME. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# GUSELKUMAB

## Products Affected

- TREMFYA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE OR GENITAL AREA.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). PSO: TRIAL OF OR CONTRAINDICATION ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. RENEWAL: PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# HIGH CONCENTRATION ORAL OPIOID SOLUTIONS

## Products Affected

- *morphine concentrate oral solution*
- *oxycodone oral concentrate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	OPIOID TOLERANT: 12 MONTHS. HOSPICE, PALLIATIVE CARE OR END OF LIFE CARE: LIFETIME.
Other Criteria	1) OPIOID TOLERANT (I.E. PREVIOUS USE OF 60 MG ORAL MORPHINE PER DAY, 25 MCG TRANSDERMAL FENTANYL PER HOUR, 30 MG ORAL OXYCODONE PER DAY, 8 MG ORAL HYDROMORPHONE PER DAY, 25 MG ORAL OXYMORPHONE PER DAY, 60 MG ORAL HYDROCODONE PER DAY, OR AN EQUIANALGESIC DOSE OF ANOTHER OPIOID) AND HAS TROUBLE SWALLOWING OPIOID TABLETS, CAPSULES, OR LARGE VOLUMES OF LIQUID, OR 2) ENROLLED IN HOSPICE OR PALLIATIVE CARE OR END OF LIFE CARE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY - ANTICHOLINERGICS - CLEMASTINE

---

## Products Affected

- *clemastine oral tablet 2.68 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# HIGH RISK DRUGS IN THE ELDERLY - ANTICHOLINERGICS - PROMETHAZINE

## Products Affected

- *promethazine injection solution*
- *promethazine oral*
- *promethazine rectal*
- *promethegan*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRURITUS/URTICARIA/SEASONAL/PERENNIAL ALLERGY: TRIAL OF OR CONTRAINDICATION TO A NON-SEDATING ANTIHISTAMINE SUCH AS LEVOCETIRIZINE OR PRESCRIBER ACKNOWLEDGEMENT OR AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. NAUSEA AND VOMITING: PRESCRIBER ACKNOWLEDGEMENT OR AWARENESS THAT THE DRUG IS CONSIDERED HIGH-RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS REQUIRE PHYSICIAN ATTESTATION THAT REQUESTED MEDICATION IS USED TO TREAT A DIAGNOSIS UNRELATED TO THE TERMINAL ILLNESS OR RELATED CONDITION, AND ARE APPROVED WITHOUT TRIAL OF FORMULARY ALTERNATIVES NOR REQUIRING PRESCRIBER ACKNOWLEDGEMENT.
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# HIGH RISK DRUGS IN THE ELDERLY - ANTICHOLINERGICS - SCOPOLAMINE

---

## Products Affected

- *scopolamine base*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS REQUIRE PHYSICIAN ATTESTATION THAT REQUESTED MEDICATION IS USED TO TREAT A DIAGNOSIS UNRELATED TO THE TERMINAL ILLNESS OR RELATED CONDITION, AND ARE APPROVED WITHOUT REQUIRING PRESCRIBER ACKNOWLEDGEMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY - BARBITURATE COMBINATIONS

## Products Affected

- *ascomp with codeine*
- *butalbital-acetaminop-caff-cod*
- *butalbital-acetaminophen oral tablet 50-325 mg*
- *butalbital-acetaminophen-caff oral capsule 50-325-40 mg*
- *butalbital-acetaminophen-caff oral tablet*
- *butalbital-aspirin-caffeine*
- *codeine-butalbital-asa-caff*
- *tencon*
- *zebutal*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# HIGH RISK DRUGS IN THE ELDERLY - DIPYRIDAMOLE

---

## Products Affected

- *dipyridamole oral*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# HIGH RISK DRUGS IN THE ELDERLY - DISOPYRAMIDE

---

## Products Affected

- *disopyramide phosphate oral capsule*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# HIGH RISK DRUGS IN THE ELDERLY - ENDOCRINE - ESTROGEN

## Products Affected

- *amabelz*
- *dotti*
- DUAVEE
- *estradiol oral*
- *estradiol transdermal patch semiweekly*
- *estradiol transdermal patch weekly*
- *estradiol-norethindrone acet oral tablet 0.5-0.1 mg*
- *fyavolv*
- *jinteli*
- *lyllana*
- *mimvey*
- *norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg*
- PREMARIN ORAL
- PREMPHASE
- PREMPRO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	VULVAR/VAGINAL ATROPHY, OSTEOPOROSIS AND VASOMOTOR SYMPTOMS OF MENOPAUSE: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. ALL OTHER FDA APPROVED INDICATIONS NOT PREVIOUSLY MENTIONED IN THIS SECTION, SUCH AS PALLIATIVE TREATMENT, AND HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# HIGH RISK DRUGS IN THE ELDERLY - ENDOCRINE - SULFONYLUREAS

---

## Products Affected

- *glyburide*
- *glyburide micronized*
- *glyburide-metformin*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF GLIMEPIRIDE, GLIPIZIDE, OR PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS ARE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No





# HIGH RISK DRUGS IN THE ELDERLY - SKELETAL MUSCLE RELAXANTS

## Products Affected

- *chlorzoxazone oral tablet 250 mg, 500 mg*
- *cyclobenzaprine oral tablet 10 mg, 5 mg*
- *methocarbamol oral tablet 500 mg, 750 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED A HIGH RISK MEDICATION FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY

## ANTICHOLINERGICS -

### CYPROHEPTADINE\_CARBINOXAMINE

---

#### Products Affected

- *carbinoxamine maleate oral liquid*
- *carbinoxamine maleate oral tablet 4 mg*
- *cyproheptadine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY- ANTICHOLINERGICS- DIPHENHYDRAMINE ELIXIR

## Products Affected

- *diphenhydramine hcl oral elixir*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ANTIHISTAMINIC CONDITIONS (PRURITUS OR URTICARIA): TRIAL OR CONTRAINDICATION TO A NON-SEDATING ANTIHISTAMINE SUCH AS LEVOCETIRIZINE. INSOMNIA: TRIAL OF SILENOR AND BELSOMRA. MOTION SICKNESS AND ANTIPARKINSONISM: PRESCRIBER ACKNOWLEDGEMENT/AWARENESS DRUG IS CONSIDERED AS HIGH RISK MEDICATION IN THE ELDERLY FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS AND ANAPHYLACTIC REACTIONS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# HIGH RISK DRUGS IN THE ELDERLY- DIPHENOXYLATE-ATROPINE

---

## Products Affected

- *diphenoxylate-atropine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY- INDOMETHACIN

---

## Products Affected

- *indomethacin oral capsule 25 mg, 50 mg*
- *indomethacin oral capsule, extended release*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY- MEGESTROL

## Products Affected

- *megestrol oral suspension 400 mg/10 ml (40 mg/ml)*
- *megestrol oral tablet*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HIGH RISK DRUGS IN THE ELDERLY- PAROXETINE

---

## Products Affected

- *paroxetine hcl*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# HIGH RISK MEDICATIONS IN THE ELDERLY- PHENOBARBITAL

---

## Products Affected

- *phenobarbital*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	APPROPRIATE HIGH RISK MEDICATION USE: 65 YEARS OF AGE OR OLDER. PA NOT REQUIRED FOR AGE 0-64 YEARS.
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	FOR TREATMENT OF EPILEPSY/SEIZURES IN PATIENTS WHO ARE NEWLY PRESCRIBED PHENOBARBITAL: PATIENT HAS NOT RESPONDED TO AT LEAST ONE ANTICONVULSANT OR PRESCRIBER ACKNOWLEDGEMENT/AWARENESS THAT THE DRUG IS CONSIDERED HIGH RISK FOR PATIENTS 65 YEARS AND OLDER. HOSPICE PATIENTS WILL BE APPROVED WITHOUT REQUIRING ADDITIONAL CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# HISTRELIN-SUPPRELIN LA

---

## Products Affected

- SUPPRELIN LA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	CENTRAL PRECOCIOUS PUBERTY (CPP): INITIAL: FEMALES: ELEVATED LEVELS OF FOLLICLE-STIMULATING HORMONE (FSH) GREATER THAN 4.0 MIU/ML AND LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVELS OF FSH GREATER THAN 5.0 MIU/ML AND LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CPP: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>CPP: INITIAL: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR BREAST DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR GENITAL DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). RENEWAL: 1) TANNER SCALE STAGING AT INITIAL DIAGNOSIS OF CPP HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) PATIENT HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# HYDROXYUREA

---

## Products Affected

- SIKLOS ORAL TABLET 100 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# IBRUTINIB

---

## Products Affected

- IMBRUVICA ORAL CAPSULE 140 MG, 70 MG
- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# IBUPROFEN-FAMOTIDINE

---

## Products Affected

- *ibuprofen-famotidine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	TRIAL OF ONE OF THE FOLLOWING GENERIC, FEDERAL LEGEND HISTAMINE H2-RECEPTOR ANTAGONISTS: FAMOTIDINE, CIMETIDINE, OR NIZATIDINE, AND TRIAL OF GENERIC, FEDERAL LEGEND IBUPROFEN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ICATIBANT

---

## Products Affected

- *icatibant*
- *sajazir*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY ANGIOEDEMA (HAE): DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
Age Restrictions	
Prescriber Restrictions	HAE: PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIST, IMMUNOLOGIST, OR HEMATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	HAE: NO CONCURRENT USE WITH OTHER MEDICATIONS FOR TREATMENT OF ACUTE HAE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# IDELALISIB

---

## Products Affected

- ZYDELIG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# IMATINIB

---

## Products Affected

- *imatinib oral tablet 100 mg, 400 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	ADJUVANT GASTROINTESTINAL STROMAL TUMOR TREATMENT: 36 MONTHS. ALL OTHER DIAGNOSES: 12 MONTHS.
Other Criteria	PHILADELPHIA CHROMOSOME POSITIVE CHRONIC MYELOID LEUKEMIA: PATIENT HAS NOT RECEIVED A PREVIOUS TREATMENT WITH ANOTHER TYROSINE KINASE INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# INFIGRATINIB

---

## Products Affected

- TRUSELTIQ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# INFLIXIMAB

## Products Affected

- *infliximab*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, SKYRIZI. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, XELJANZ, RINVOQ. RENEWAL: RA, AS, PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# INFLIXIMAB-ABDA

---

## Products Affected

- RENFLEXIS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, SKYRIZI. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, XELJANZ, RINVOQ. RENEWAL: RA, AS, PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# INFLIXIMAB-AXXQ

---

## Products Affected

- AVSOLA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, SKYRIZI. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, XELJANZ, RINVOQ. RENEWAL: RA, AS, PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# INFLIXIMAB-DYYB

## Products Affected

- INFLECTRA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, COSENTYX, ENBREL, SKYRIZI, TREMFYA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, ENBREL, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, SKYRIZI. UC: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, STELARA, XELJANZ, RINVOQ. RENEWAL: RA, AS, PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# INTERFERON ALFA-2B

## Products Affected

- INTRON A INJECTION RECON SOLN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HEPATITIS C: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (E.G. HEPATOLOGIST).
Coverage Duration	6 MONTHS.
Other Criteria	HEPATITIS C: 1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) TRIAL OF OR CONTRAINDICATION TO PEGINTERFERON ALFA-2A OR PEGINTERFERON ALFA-2B, 3) USED IN COMBINATION WITH RIBAVIRIN UNLESS CONTRAINDICATED AND 4) LIMITED TO TOTAL OF 24 MONTHS OF TREATMENT. FOLLICULAR LYMPHOMA: LIMITED TO TOTAL OF 18 MONTHS OF TREATMENT. ALL OTHER INDICATIONS: LIMITED TO TOTAL OF 1 YEAR OF TREATMENT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# INTERFERON FOR MS-AVONEX

---

## Products Affected

- AVONEX INTRAMUSCULAR PEN INJECTOR KIT
- AVONEX INTRAMUSCULAR SYRINGE KIT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# INTERFERON FOR MS-BETASERON

---

## Products Affected

- BETASERON SUBCUTANEOUS KIT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# INTERFERON FOR MS-PLEGRIDY

---

## Products Affected

- PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML
- PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# INTERFERON GAMMA-1B

## Products Affected

- ACTIMMUNE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: CHRONIC GRANULOMATOUS DISEASE (CGD): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, INFECTIOUS DISEASE SPECIALIST, OR IMMUNOLOGIST. SEVERE MALIGNANT OSTEOPETROSIS (SMO): PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	RENEWAL: CGD, SMO: 1) DEMONSTRATED CLINICAL BENEFIT COMPARED TO BASELINE, AND 2) HAS NOT RECEIVED HEMATOPOIETIC CELL TRANSPLANTATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# IPILIMUMAB

---

## Products Affected

- YERVOY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: UNRESECT/MET MEL: 4MO, RCC/CRC/HCC: 3MO, ALL OTHERS: 12MO. INITIAL/RENEWAL: CUTAN MEL: 6MO
Other Criteria	RENEWAL: ADJUVANT CUTANEOUS MELANOMA: NO EVIDENCE OF DISEASE RECURRENCE (DEFINED AS THE APPEARANCE OF ONE OR MORE NEW MELANOMA LESIONS: LOCAL, REGIONAL OR DISTANT METASTASIS). THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# ITRACONAZOLE SOLUTION

---

## Products Affected

- *itraconazole oral solution*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS
Other Criteria	ESOPHAGEAL CANDIDIASIS AND OROPHARYNGEAL CANDIDIASIS; TRIAL OF OR CONTRAINDICATION TO FLUCONAZOLE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# IVACAF TOR

---

## Products Affected

- KALYDECO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CYSTIC FIBROSIS (CF); INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: LIFETIME
Other Criteria	CF: INITIAL: NOT HOMOZYGOUS FOR F508DEL MUTATION IN CFTR GENE. RENEWAL: 1) MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR 2) REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# IVOSIDENIB

---

## Products Affected

- TIBSOVO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# IXAZOMIB

---

## Products Affected

- NINLARO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# IXEKIZUMAB

## Products Affected

- TALTZ AUTOINJECTOR
- TALTZ SYRINGE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, OR GENITAL AREA. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: HUMIRA, COSENTYX, STELARA, ENBREL, SKYRIZI, TREMFYA. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, STELARA, ENBREL, XELJANZ, TREMFYA, RINVOQ, SKYRIZI. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA, COSENTYX, XELJANZ, RINVOQ. NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, RINVOQ. RENEWAL: PSO, PSA, AS, NR-AXSPA: CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LANADELUMAB

---

## Products Affected

- TAKHZYRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.
Age Restrictions	
Prescriber Restrictions	HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIST, IMMUNOLOGIST, OR HEMATOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT COMPARED TO BASELINE IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LANREOTIDE

## Products Affected

- *lanreotide* ML
- SOMATULINE DEPOT  
SUBCUTANEOUS SYRINGE 120  
MG/0.5 ML, 60 MG/0.2 ML, 90 MG/0.3

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	ACROMEGALY: INITIAL: THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	ACROMEGALY: INITIAL: 3 MOS, RENEWAL: 12 MOS.GEP-NETS, CARCINOID SYNDROME: 12 MOS.
Other Criteria	ACROMEGALY: INITIAL: TRIAL OF OR CONTRAINDICATION TO ONE GENERIC OCTREOTIDE INJECTION. RENEWAL: 1) REDUCTION, NORMALIZATION, OR MAINTENANCE OF IGF-1 LEVELS BASED ON AGE AND GENDER, AND 2) IMPROVEMENT OR SUSTAINED REMISSION OF CLINICAL SYMPTOMS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# LAPATINIB

---

## Products Affected

- *lapatinib*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LAROTRECTINIB

---

## Products Affected

- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	APPROVAL FOR VITRAKVI ORAL SOLUTION: TRIAL OF VITRAKVI CAPSULES OR PATIENT IS UNABLE TO TAKE CAPSULE FORMULATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LEDIPASVIR-SOFOSBUVIR

## Products Affected

- HARVONI ORAL PELLETS IN PACKET 33.75-150 MG, 45-200 MG
- HARVONI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, AND 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING: CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, ROSUVASTATIN, TIPRANA VIR/RITONAVIR, SOFOSBUVIR (AS A SINGLE AGENT), EPCLUSA, ZEPATIER, MAVYRET, OR VOSEVI. REQUESTS FOR HARVONI 45MG-200MG PELLETS: PATIENT IS UNABLE TO SWALLOW TABLETS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LENALIDOMIDE

---

## Products Affected

- *lenalidomide*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LENVATINIB

---

## Products Affected

- LENVIMA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LETERMIVIR

---

## Products Affected

- PREVYMIS INTRAVENOUS SOLUTION 240 MG/12 ML, 480 MG/24 ML
- PREVYMIS ORAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	4 MONTHS
Other Criteria	HEMATOPOIETIC STEM CELL TRANSPLANT (HSCT): THERAPY WILL BE INITIATED BETWEEN DAY 0 AND DAY 28 POST TRANSPLANTATION AND PATIENT IS NOT RECEIVING THE MEDICATION BEYOND 100 DAYS POST TRANSPLANTATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LEUPROLIDE

## Products Affected

- *leuprolide subcutaneous kit*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: CENTRAL PRECOCIOUS PUBERTY (CPP): FEMALES: ELEVATED LEVELS OF FOLLICLE-STIMULATING HORMONE (FSH) GREATER THAN 4.0 MIU/ML AND LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVELS OF FSH GREATER THAN 5.0 MIU/ML AND LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: CPP: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	PROSTATE CANCER: 12 MONTHS. CPP: INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	CPP: INITIAL: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR BREAST DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR GENITAL DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). RENEWAL: 1) TANNER SCALE STAGING AT INITIAL DIAGNOSIS OF CPP HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) PATIENT HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# LEUPROLIDE DEPOT

---

## Products Affected

- *leuprolide (3 month)*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LEUPROLIDE-ELIGARD

---

## Products Affected

- ELIGARD
- ELIGARD (3 MONTH)
- ELIGARD (4 MONTH)
- ELIGARD (6 MONTH)

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS.
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LEUPROLIDE-LUPRON DEPOT

## Products Affected

- LUPRON DEPOT
- LUPRON DEPOT (3 MONTH)
- LUPRON DEPOT (4 MONTH)
- LUPRON DEPOT (6 MONTH)

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.
<b>Coverage Duration</b>	PROSTATE CA: 12 MOS. UTERINE FIBROIDS: 3 MOS. ENDOMETRIOSIS: INITIAL/RENEWAL: 6 MOS.
<b>Other Criteria</b>	INITIAL: ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME. RENEWAL: ENDOMETRIOSIS: 1) IMPROVEMENT OF PAIN RELATED TO ENDOMETRIOSIS WHILE ON THERAPY, 2) RECEIVING CONCOMITANT ADD-BACK THERAPY (I.E., COMBINATION ESTROGEN-PROGESTIN OR PROGESTIN-ONLY CONTRACEPTIVE PREPARATION), 3) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 4) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LEUPROLIDE-LUPRON DEPOT-PED

## Products Affected

- LUPRON DEPOT-PED (3 MONTH) INTRAMUSCULAR SYRINGE KIT 30 MG
- LUPRON DEPOT-PED INTRAMUSCULAR KIT 11.25 MG, 15 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	CENTRAL PRECOCIOUS PUBERTY (CPP): INITIAL: FEMALES: ELEVATED LEVELS OF FOLLICLE-STIMULATING HORMONE (FSH) GREATER THAN 4.0 MIU/ML AND LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVELS OF FSH GREATER THAN 5.0 MIU/ML AND LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS.
Age Restrictions	
Prescriber Restrictions	CPP: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	<p>CPP: INITIAL: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR BREAST DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR GENITAL DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). RENEWAL: 1) TANNER SCALE STAGING AT INITIAL DIAGNOSIS OF CPP HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) PATIENT HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LEVODOPA

---

## Products Affected

- INBRIJA INHALATION CAPSULE,  
W/INHALATION DEVICE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	PD: INITIAL: 1) NOT CURRENTLY TAKING MORE THAN 1600MG OF LEVODOPA PER DAY, AND 2) PHYSICIAN HAS OPTIMIZED DRUG THERAPY FOR PARKINSONS DISEASE. RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WITH THE USE OF INBRIJA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# L-GLUTAMINE

---

## Products Affected

- ENDARI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SICKLE CELL DISEASE(SCD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST
Coverage Duration	INITIAL: 12 MONTHS. RENEWAL: LIFETIME.
Other Criteria	SCD: INITIAL: PATIENTS 18 YEARS OR OLDER: ONE OF THE FOLLOWING: 1) AT LEAST 2 SICKLE CELL CRISES IN THE PAST YEAR, 2) SICKLE-CELL ASSOCIATED SYMPTOMS WHICH ARE INTERFERING WITH ACTIVITIES OF DAILY LIVING, OR 3) HISTORY OF OR HAS RECURRENT ACUTE CHEST SYNDROME. PATIENTS 5 TO 17 YEARS: APPROVED WITHOUT ADDITIONAL CRITERIA. RENEWAL: HAS MAINTAINED OR EXPERIENCED REDUCTION IN ACUTE COMPLICATIONS OF SICKLE CELL DISEASE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# LIDOCAINE

## Products Affected

- *lidocaine hcl mucous membrane solution 4 % (40 mg/ml)*
- *lidocaine topical adhesive patch, medicated 5 %*
- *lidocaine topical ointment*
- **ZTLIDO**

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
<b>Indications</b>	All Medically-accepted Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LIDOCAINE PRILOCAINE

---

## Products Affected

- *lidocaine-prilocaine topical cream*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# LOMITAPIDE

---

## Products Affected

- JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 30 MG, 40 MG, 5 MG, 60 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST, ENDOCRINOLOGIST OR LIPIDOLOGIST.
<b>Coverage Duration</b>	12 MONTHS

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>1) DIAGNOSIS DETERMINED BY A) DEFINITE SIMON BROOME DIAGNOSTIC CRITERIA, OR B) DUTCH LIPID NETWORK CRITERIA SCORE OF 8 OR GREATER, OR C) CLINICAL DIAGNOSIS BASED ON A HISTORY OF AN UNTREATED LDL-C CONCENTRATION GREATER THAN 500 MG/DL TOGETHER WITH EITHER XANTHOMA BEFORE 10 YEARS OF AGE, OR EVIDENCE OF HEFH IN BOTH PARENTS. 2) LDL-C LEVEL GREATER THAN OR EQUAL TO 70MG/DL WHILE ON MAXIMAL DRUG TREATMENT. 3) TRIAL OF EVOLOCUMAB UNLESS THE PATIENT HAS NON-FUNCTIONING LDL RECEPTORS. 4) MEETS ONE OF THE FOLLOWING: A) TAKING A HIGH-INTENSITY STATIN (I.E., ATORVASTATIN 40-80MG DAILY, ROSUVASTATIN 20-40MG DAILY) FOR A DURATION OF AT LEAST 8 WEEKS, B) TAKING A MAXIMALLY TOLERATED DOSE OF ANY STATIN FOR A DURATION OF AT LEAST 8 WEEKS GIVEN THAT THE PATIENT CANNOT TOLERATE A HIGH-INTENSITY STATIN, C) ABSOLUTE CONTRAINDICATION TO STATIN THERAPY (E.G., ACTIVE DECOMPENSATED LIVER DISEASE, NURSING FEMALE, PREGNANCY OR PLANS TO BECOME PREGNANT, HYPERSENSITIVITY REACTIONS), D) STATIN INTOLERANCE, OR E) TRIAL OF ROSUVASTATIN, ATORVASTATIN, OR STATIN THERAPY AT ANY DOSE AND HAS EXPERIENCED SKELETAL-MUSCLE RELATED SYMPTOMS (E.G., MYOPATHY).</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LONCASTUXIMAB TESIRINE-LPYL

---

## Products Affected

- ZYNLONTA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LORLATINIB

---

## Products Affected

- LORBRENA ORAL TABLET 100 MG,  
25 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# LUMACAF TOR-IVACAF TOR

## Products Affected

- ORKAMBI ORAL GRANULES IN PACKET
- ORKAMBI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	INITIAL: CYSTIC FIBROSIS (CF): CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CF.
Age Restrictions	
Prescriber Restrictions	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CF EXPERT.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: LIFETIME.
Other Criteria	CF: RENEWAL: MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# LUMASIRAN

---

## Products Affected

- OXLUMO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# MACITENTAN

## Products Affected

- OPSUMIT

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: 1) DOCUMENTED CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) OF 3 WOOD UNITS OR GREATER, AND 2) NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PAH: RENEWAL: 1) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE, OR 2) A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MARGETUXIMAB-CMKB

---

## Products Affected

- MARGENZA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MEPOLIZUMAB

## Products Affected

- NUCALA SUBCUTANEOUS AUTO-INJECTOR
- NUCALA SUBCUTANEOUS RECON SOLN
- NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: ASTHMA: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN PULMONARY OR ALLERGY MEDICINE. NASAL POLYPS: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST.
<b>Coverage Duration</b>	INITIAL: ASTHMA: 4 MO. NASAL POLYPS: 6 MO. OTHERS: 12 MO. RENEWAL: NASAL POLYPS, ASTHMA: 12 MO.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE OF THE FOLLOWING: (A) PATIENT EXPERIENCED AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) PATIENT HAS POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA. NASAL POLYPS: PREVIOUS 90 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID. RENEWAL: ASTHMA: 1) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR D) INCREASE IN PERCENT PREDICTED FEV1 FROM</p>
	<p>PRETREATMENT BASELINE. NASAL POLYPS: CLINICAL BENEFIT COMPARED TO BASELINE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# METHYLNALTREXONE

## Products Affected

- RELISTOR SUBCUTANEOUS SOLUTION
- RELISTOR SUBCUTANEOUS SYRINGE 12 MG/0.6 ML, 8 MG/0.4 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	ADVANCED ILLNESS: OPIOID-INDUCED CONSTIPATION.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 MONTHS FOR PATIENTS RECEIVING PALLIATIVE CARE, 12 MONTHS FOR CHRONIC, NON-CANCER PAIN.
Other Criteria	ADVANCED ILLNESS: PATIENT IS RECEIVING PALLIATIVE CARE. CHRONIC NON-CANCER PAIN: PATIENT HAS BEEN TAKING OPIOIDS FOR AT LEAST 4 WEEKS AND HAD A PREVIOUS TRIAL OF OR CONTRAINDICATION TO NALOXEGOL (MOVANTIK) AND LUBIPROSTONE (AMITIZA).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# METHYLNALTREXONE ORAL

---

## Products Affected

- RELISTOR ORAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	OPIOID INDUCED CONSTIPATION WITH CHRONIC NON-CANCER PAIN: 1) HAS BEEN TAKING OPIOIDS FOR AT LEAST 4 WEEKS, AND 2) TRIAL OF OR CONTRAINDICATION TO NALOXEGOL (MOVANTIK) AND LUBIPROSTONE (AMITIZA).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# MIDOSTAURIN

---

## Products Affected

- RYDAPT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	ACUTE MYELOID LEUKEMIA: 6 MONTHS. ADVANCED SYSTEMIC MASTOCYTOSIS: 12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# MIFEPRISTONE

## Products Affected

- KORLYM

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	CUSHINGS SYNDROME (CS): INITIAL: DIAGNOSIS CONFIRMED BY ONE OF THE FOLLOWING: 1) 24-HR URINE FREE CORTISOL (2 OR MORE TESTS TO CONFIRM), 2) OVERNIGHT 1MG DEXAMETHASONE TEST, OR 3) LATE NIGHT SALIVARY CORTISOL (2 OR MORE TESTS TO CONFIRM).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CS: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	INITIAL AND RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	CS: INITIAL: HYPERCORTISOLISM IS NOT A RESULT OF CHRONIC GLUCOCORTICOIDS. RENEWAL: 1) CONTINUES TO HAVE IMPROVEMENT OF GLUCOSE TOLERANCE OR STABLE GLUCOSE TOLERANCE (E.G., REDUCED A1C, IMPROVED FASTING GLUCOSE, ETC.), 2) CONTINUES TO HAVE TOLERABILITY TO KORLYM, AND 3) CONTINUES TO NOT BE A CANDIDATE FOR SURGICAL TREATMENT OR HAS FAILED SURGERY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MIGALASTAT

## Products Affected

- GALAFOLD

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	FABRY DISEASE: INITIAL: PATIENT IS SYMPTOMATIC OR HAS EVIDENCE OF INJURY FROM GL-3 TO THE KIDNEY, HEART, OR CENTRAL NERVOUS SYSTEM RECOGNIZED BY LABORATORY, HISTOLOGICAL, OR IMAGING FINDINGS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	FABRY DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST, CARDIOLOGIST, OR SPECIALIST IN GENETICS OR INHERITED METABOLIC DISORDERS.
<b>Coverage Duration</b>	INITIAL: 6 MOS. RENEWAL: 12 MOS.
<b>Other Criteria</b>	FABRY DISEASE: INITIAL: NOT CONCURRENTLY USING ENZYME REPLACEMENT THERAPY (I.E. FABRAZYME), RENEWAL: 1) PATIENT HAS DEMONSTRATED IMPROVEMENT OR STABILIZATION, AND 2) NOT CONCURRENTLY USING ENZYME REPLACEMENT THERAPY (I.E. FABRAZYME).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MIGLUSTAT

---

## Products Affected

- *miglustat*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MILTEFOSINE

---

## Products Affected

- IMPAVIDO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MOBOCERTINIB

---

## Products Affected

- EXKIVITY

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# MOSUNETUZUMAB-AXGB

---

## Products Affected

- LUNSUMIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	RELAPSED OR REFRACTORY FOLLICULAR LYMPHOMA: INITIAL: 6 MONTHS. RENEWAL: 7 MONTHS.
Other Criteria	RELAPSED OR REFRACTORY FOLLICULAR LYMPHOMA: RENEWAL: 1) HAS ACHIEVED A PARTIAL RESPONSE TO TREATMENT, AND 2) HAS NOT PREVIOUSLY RECEIVED MORE THAN 17 CYCLES OF TREATMENT. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# NAFARELIN

---

## Products Affected

- SYNAREL

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS. CENTRAL PRECOCIOUS PUBERTY (CPP): FEMALES: ELEVATED LEVELS OF FOLLICLE-STIMULATING HORMONE (FSH) GREATER THAN 4.0 MIU/ML AND LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVELS OF FSH GREATER THAN 5.0 MIU/ML AND LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST. CPP: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	ENDOMETRIOSIS: 6 MONTHS. CPP: INITIAL/RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 6 MONTHS OF TREATMENT PER LIFETIME. CPP: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR BREAST DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR GENITAL DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). RENEWAL: CPP: 1) TANNER SCALE STAGING AT INITIAL DIAGNOSIS OF CPP HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# NARCOLEPSY AGENTS

---

## Products Affected

- *armodafinil*
- *modafinil oral tablet 100 mg, 200 mg*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# NATALIZUMAB

## Products Affected

- TYSABRI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	MULTIPLE SCLEROSIS (MS): 12 MOS. CD: INITIAL: 6 MOS, RENEWAL: 12 MOS.
<b>Other Criteria</b>	INITIAL: MS: TRIAL OF TWO AGENTS INDICATED FOR THE TREATMENT OF MS. CD: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, STELARA, SKYRIZI. RENEWAL: CD: ONE OF THE FOLLOWING: 1) RECEIVED AT LEAST 12 MONTHS OF THERAPY WITH TYSABRI AND HAS NOT REQUIRED MORE THAN 3 MONTHS OF CORTICOSTEROID USE WITHIN THE PAST 12 MONTHS TO CONTROL THEIR CROHNS DISEASE WHILE ON TYSABRI, OR 2) HAS ONLY RECEIVED 6 MONTHS OF THERAPY WITH TYSABRI AND HAS TAPERED OFF CORTICOSTEROIDS DURING THE FIRST 24 WEEKS OF TYSABRI THERAPY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# NAXITAMAB-GQGK

---

## Products Affected

- DANYELZA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# NERATINIB MALEATE

---

## Products Affected

- NERLYNX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	EARLY-STAGE (STAGE I-III) BREAST CANCER: MEDICATION IS BEING REQUESTED WITHIN 2 YEARS OF COMPLETING THE LAST TRASTUZUMAB DOSE. ALL OTHER FDA APPROVED INDICATIONS ARE COVERED WITHOUT ADDITIONAL CRITERIA, EXCEPT THOSE CRITERIA IN THE FDA APPROVED LABEL.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# NILOTINIB

---

## Products Affected

- TASIGNA ORAL CAPSULE 150 MG, 200 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PREVIOUSLY TREATED CML: MUTATIONAL ANALYSIS PRIOR TO INITIATION AND TASIGNA IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# NINTEDANIB

## Products Affected

- OFEV

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	<p>INITIAL: IDIOPATHIC PULMONARY FIBROSIS (IPF): 1) A USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT, AND 2) BASELINE FORCED VITAL CAPACITY (FVC) AT LEAST 50% OF PREDICTED VALUE. SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE (SSC-ILD): 1) AT LEAST 10% FIBROSIS ON A CHEST HRCT, AND 2) BASELINE FVC AT LEAST 40% OF PREDICTED VALUE. CHRONIC FIBROSING INTERSTITIAL LUNG DISEASE WITH A PROGRESSIVE PHENOTYPE (PF-ILD): 1) AT LEAST 10% FIBROSIS ON A CHEST HRCT, AND 2) BASELINE FVC AT LEAST 45% OF PREDICTED VALUE.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	<p>INITIAL: IPF: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST. SSC-ILD, PF-ILD: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR RHEUMATOLOGIST.</p>
<b>Coverage Duration</b>	<p>INITIAL: SSC-ILD: 6 MOS. IPF, PF-ILD: 12 MOS. RENEWAL (ALL INDICATIONS): 12 MOS.</p>

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: IPF: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS), AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: ESBRIET (PIRFENIDONE). SSC-ILD: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., HEART FAILURE/FLUID OVERLOAD, DRUG-INDUCED LUNG TOXICITY, RECURRENT ASPIRATION), AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: ACTEMRA SUBQ. PF-ILD: LUNG FUNCTION AND RESPIRATORY SYMPTOMS OR CHEST IMAGING HAVE WORSENERD/PROGRESSED DESPITE TREATMENT WITH MEDICATIONS USED IN CLINICAL PRACTICE FOR ILD (NOT ATTRIBUTABLE TO COMORBIDITIES SUCH AS INFECTION, HEART FAILURE). RENEWAL: IPF, SSC-ILD, PF-ILD: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# NIRAPARIB

---

## Products Affected

- ZEJULA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	RECURRENT EPITHELIAL OVARIAN, FALLOPIAN TUBE, OR PRIMARY PERITONEAL CANCER: 1) ZEJULA WILL BE USED AS MONOTHERAPY, AND 2) ZEJULA IS STARTED NO LATER THAN 8 WEEKS AFTER THE MOST RECENT PLATINUM-CONTAINING REGIMEN.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# NITISINONE

## Products Affected

- *nitisinone*
- ORFADIN ORAL CAPSULE 20 MG
- ORFADIN ORAL SUSPENSION

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HEREDITARY TYROSINEMIA TYPE 1 (HT-1): INITIAL: DIAGNOSIS CONFIRMED BY ELEVATED URINARY OR PLASMA SUCCINYLACETONE LEVELS OR A MUTATION IN THE FUMARYLACETOACETATE HYDROLASE GENE. RENEWAL: URINARY OR PLASMA SUCCINYLACETONE LEVELS HAVE DECREASED FROM BASELINE WHILE ON TREATMENT WITH NITISINONE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HT-1: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PRESCRIBER SPECIALIZING IN INHERITED METABOLIC DISEASES.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	HT-1: INITIAL: ORFADIN SUSPENSION: TRIAL OF OR CONTRAINDICATION TO PREFERRED NITISINONE TABLETS OR CAPSULES.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# NIVOLUMAB

---

## Products Affected

- OPDIVO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	UNRESECTABLE OR METASTATIC MELANOMA: NO CONCURRENT USE WITH TARGETED THERAPY (I.E., BRAF INHIBITORS, MEK INHIBITORS, AND NTRK INHIBITORS).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# NIVOLUMAB-RELATLIMAB-RMBW

---

## Products Affected

- OPDUALAG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OBETICHOLIC ACID

## Products Affected

- OCALIVA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	PRIMARY BILIARY CHOLANGITIS (PBC): INITIAL/RENEWAL: COMPLETE BILIARY OBSTRUCTION.
<b>Required Medical Information</b>	PBC: INITIAL: DIAGNOSIS CONFIRMED BY TWO OF THE FOLLOWING: 1) ALKALINE PHOSPHATASE LEVEL OF AT LEAST 1.5 TIMES THE UPPER LIMIT OF NORMAL, 2) PRESENCE OF ANTIMITOCHONDRIAL ANTIBODIES AT A TITER OF 1:40 OR HIGHER, OR 3) HISTOLOGIC EVIDENCE OF NON-SUPPURATIVE DESTRUCTIVE CHOLANGITIS AND DESTRUCTION OF INTERLOBULAR BILE DUCTS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PBC: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST OR HEPATOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	PBC: INITIAL: USED IN COMBINATION WITH URSODEOXYCHOLIC ACID IN A PATIENT WITH AN INADEQUATE RESPONSE TO URSODEOXYCHOLIC ACID, OR AS MONOTHERAPY IN A PATIENT WHO IS UNABLE TO TOLERATE URSODEOXYCHOLIC ACID. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OCRELIZUMAB

---

## Products Affected

- OCREVUS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	RELAPSING FORM OF MULTIPLE SCLEROSIS (MS): TRIAL OF TWO AGENTS INDICATED FOR THE TREATMENT OF RELAPSING FORMS OF MS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OFATUMUMAB-SQ

---

## Products Affected

- KESIMPTA PEN

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OLANZAPINE/SAMIDORPHAN

## Products Affected

- LYBALVI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SCHIZOPHRENIA/BIPOLAR I: PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST
Coverage Duration	12 MONTHS
Other Criteria	SCHIZOPHRENIA: (1) PATIENT IS AT HIGH RISK OF WEIGHT GAIN AND (2) TRIAL OF OR CONTRAINDICATION TO LATUDA OR ONE OF THE FOLLOWING ORAL ANTIPSYCHOTICS: RISPERIDONE, CLOZAPINE TABLET, OLANZAPINE, IMMEDIATE RELEASE QUETIAPINE FUMARATE, ZIPRASIDONE, ARIPIPRAZOLE. BIPOLAR I: (1) PATIENT IS AT HIGH RISK OF WEIGHT GAIN AND (2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING ORAL ANTIPSYCHOTICS: RISPERIDONE, OLANZAPINE, IMMEDIATE RELEASE QUETIAPINE FUMARATE, ZIPRASIDONE, ARIPIPRAZOLE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# OLAPARIB

## Products Affected

- LYNPARZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	RECURRENT EPITHELIAL OVARIAN, FALLOPIAN TUBE OR PRIMARY PERITONEAL CANCER: 1) MEDICATION WILL BE USED AS MONOTHERAPY, AND 2) PATIENT HAS COMPLETED TWO OR MORE LINES OF PLATINUM-BASED CHEMOTHERAPY. METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: 1) PREVIOUSLY RECEIVED A BILATERAL ORCHIECTOMY, OR 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. ALL OTHER FDA APPROVED INDICATIONS ARE COVERED WITHOUT ADDITIONAL CRITERIA, EXCEPT THOSE CRITERIA IN THE FDA APPROVED LABEL.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# OLUTASIDENIB

---

## Products Affected

- REZLIDHIA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OMACETAXINE

---

## Products Affected

- SYNRIBO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OMALIZUMAB

---

## Products Affected

- XOLAIR

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: ASTHMA: POSITIVE SKIN PRICK OR BLOOD TEST (E.G., ELISA, FEIA) TO A PERENNIAL AEROALLERGEN AND A BASELINE IGE SERUM LEVEL GREATER THAN OR EQUAL TO 30 IU/ML.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL AND RENEWAL: CHRONIC IDIOPATHIC URTICARIA (CIU): PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE, DERMATOLOGY OR IMMUNOLOGY. INITIAL: NASAL POLYPS: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.
<b>Coverage Duration</b>	INITIAL: ASTHMA: 12 MO. CIU, NASAL POLYPS: 6 MO. RENEWAL: ASTHMA, NASAL POLYPS: 12 MO. CIU: 6 MO.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: CIU: TRIAL OF OR CONTRAINDICATION TO A MAXIMALLY TOLERATED DOSE OF AN H1 ANTI-HISTAMINE AND STILL EXPERIENCES HIVES ON MOST DAYS OF THE WEEK. NASAL POLYPS: 1) PREVIOUS 90 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE PREFERRED AGENT: NUCALA, DUPIXENT. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, 2) ONE OF THE FOLLOWING: (A) PATIENT EXPERIENCED AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) PATIENT HAS POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, 3) NOT CONCURRENTLY RECEIVING DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA. RENEWAL: CIU: DIAGNOSIS OF CIU. NASAL POLYPS: CLINICAL BENEFIT COMPARED TO BASELINE. ASTHMA: 1) NOT CONCURRENTLY RECEIVING DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: A) REDUCTION IN ASTHMA</p>
	<p>EXACERBATIONS FROM BASELINE, B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR D) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE.</p>
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OPICAPONE

---

## Products Affected

- ONGENTYS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	PARKINSONS DISEASE: 18 YEARS OF AGE OR OLDER
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# OSIMERTINIB

---

## Products Affected

- TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	EGFR EXON 19 DELETIONS OR EXON 21 L858R MUTATIONS NON-SMALL CELL LUNG CANCER (NSCLC) AND METASTATIC NSCLC WITH EGFR T790M MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PACRITINIB

---

## Products Affected

- VONJO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MYELOFIBROSIS: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# PALBOCICLIB

---

## Products Affected

- IBRANCE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PALIVIZUMAB

## Products Affected

- SYNAGIS

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: GESTATIONAL AGE
<b>Age Restrictions</b>	INITIAL AND RENEWAL: LESS THAN 24 MONTHS OF AGE.
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	CRITERIA CONSISTENT WITH THE CDC RSV CENSUS REGIONAL TREND. SEE OTHER CRITERIA.
<b>Other Criteria</b>	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT RECOMMENDATIONS FROM THE AMERICAN ACADEMY OF PEDIATRICS FOR PALIVIZUMAB PROPHYLAXIS FOR RESPIRATORY SYNCYTIAL VIRUS (RSV) INFECTIONS AND THE CENTERS FOR DISEASE CONTROL AND PREVENTION RSV CENSUS REGIONAL TREND. APPROVAL WILL BE FOR UP TO A MAXIMUM OF 5 MONTHS DEPENDENT UPON REMAINING LENGTH OF RESPIRATORY SYNCYTIAL VIRUS SEASON. RENEWAL: ADDITIONAL 1 MONTH OF TREATMENT FOR CARDIOPULMONARY BYPASS SURGERY DURING RSV PROPHYLAXIS SEASON.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PANOBINOSTAT

---

## Products Affected

- FARYDAK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	MULTIPLE MYELOMA: RENEWAL: TOLERATED THE FIRST 8 CYCLES OF THERAPY WITHOUT UNRESOLVED SEVERE OR MEDICALLY SIGNIFICANT TOXICITY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PARATHYROID HORMONE

---

## Products Affected

- NATPARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HYPOCALCEMIA SECONDARY TO HYPOPARATHYROIDISM: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	HYPOCALCEMIA SECONDARY TO HYPOPARATHYROIDISM: 1) TRIAL OF OR CONTRAINDICATION TO CALCITRIOL, 2) HYPOPARATHYROIDISM IS NOT DUE TO A CALCIUM SENSING RECEPTOR (CSR) MUTATION, AND 3) HYPOPARATHYROIDISM IS NOT CONSIDERED ACUTE POST-SURGICAL HYPOPARATHYROIDISM.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PASIREOTIDE DIASPARTATE

---

## Products Affected

- SIGNIFOR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CUSHINGS DISEASE (CD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	CD: RENEWAL: 1) CONTINUED IMPROVEMENT OF CUSHINGS DISEASE, AND 2) MAINTAINED TOLERABILITY TO SIGNIFOR.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PAZOPANIB

---

## Products Affected

- VOTRIENT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	ADVANCED SOFT TISSUE SARCOMA (STS): NOT USED FOR ADIPOCYTIC STS OR GASTROINTESTINAL STROMAL TUMORS (GIST)
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PDE5 INHIBITORS FOR PULMONARY ARTERIAL HYPERTENSION

## Products Affected

- *alyq*
- *sildenafil (pulm.hypertension) oral tablet*
- *tadalafil (pulm. hypertension)*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	DOCUMENTED CONFIRMATORY PULMONARY ARTERIAL HYPERTENSION (PAH) DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION. PATIENT HAS NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	PAH: INITIAL: 1) NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA), ANY ORGANIC NITRATES IN ANY FORM, OR GUANYLATE CYCLASE STIMULATORS, AND 2) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) 15 MMHG OR LESS, PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER. RENEWAL: 1) NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA), ANY ORGANIC NITRATES IN ANY FORM, OR GUANYLATE CYCLASE STIMULATORS, AND 2) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# PDE5 INHIBITORS FOR PULMONARY ARTERIAL HYPERTENSION - IV

## Products Affected

- *sildenafil (pulm.hypertension) intravenous*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	DOCUMENTED CONFIRMATORY PULMONARY ARTERIAL HYPERTENSION (PAH) DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION. PATIENT HAS NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL AND RENEWAL: 12 MONTHS
<b>Other Criteria</b>	PAH: INITIAL: 1) NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA), ANY ORGANIC NITRATES IN ANY FORM, OR GUANYLATE CYCLASE STIMULATORS, AND 2) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) 15 MMHG OR LESS, PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER, RENEWAL: 1) NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA), ANY ORGANIC NITRATES IN ANY FORM, OR GUANYLATE CYCLASE STIMULATORS, AND 2) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEGFILGRASTIM

---

## Products Affected

- NEULASTA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NON MYELOID MALIGNANCY, ACUTE RADIATION EXPOSURE: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEGFILGRASTIM - APGF

---

## Products Affected

- NYVEPRIA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEGFILGRASTIM - CBQV

---

## Products Affected

- UDENYCA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEGFILGRASTIM - JMDB

---

## Products Affected

- FULPHILA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
Coverage Duration	12 MONTHS
Other Criteria	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# PEGFILGRASTIM-BMEZ

---

## Products Affected

- ZIEXTENZO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO NYVEPRIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEGFILGRASTIM-NEULASTA ONPRO

---

## Products Affected

- NEULASTA ONPRO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NON MYELOID MALIGNANCY, ACUTE RADIATION EXPOSURE: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# PEGFILGRASTIM-PBBK

---

## Products Affected

- FYLNETRA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NON MYELOID MALIGNANCY: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	NON MYELOID MALIGNANCY: TRIAL OF OR CONTRAINDICATION TO PREFERRED AGENT: NYVEPRIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEGVALIASE-PQPZ

---

## Products Affected

- PALYNZIQ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PHENYLKETONURIA (PKU): INITIAL: NOT ON CONCURRENT TREATMENT WITH KUVAN. RENEWAL: 1) PATIENT CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NOT ON CONCURRENT TREATMENT WITH KUVAN .
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEGVISOMANT

---

## Products Affected

- SOMAVERT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEMBROLIZUMAB

---

## Products Affected

- KEYTRUDA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	UNRESECTABLE OR METASTATIC MELANOMA: NO CONCURRENT USE WITH TARGETED THERAPY (I.E., BRAF INHIBITORS, MEK INHIBITORS, AND NTRK INHIBITORS).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEMIGATINIB

---

## Products Affected

- PEMAZYRE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PENICILLAMINE

## Products Affected

- *penicillamine oral tablet*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	<p>INITIAL: WILSONS DISEASE: CONFIRMED BY ONE OF THE FOLLOWING: 1) PLASMA COPPER-PROTEIN CERULOPLASMIN IS LESS THAN 20MG/DL, 2) LIVER BIOPSY POSITIVE FOR AN ABNORMALLY HIGH CONCENTRATION OF COPPER (GREATER THAN 250MCG/G DRY WEIGHT) OR THE PRESENCE OF KAYSER-FLEISCHER RINGS, OR 3) CONFIRMATION BY GENETIC TESTING FOR ATP7B MUTATIONS. CYSTINURIA: PATIENT HAS NEPHROLITHIASIS AND ONE OR MORE OF THE FOLLOWING: 1) STONE ANALYSIS SHOWING PRESENCE OF CYSTEINE, 2) IDENTIFICATION OF PATHOGNOMONIC HEXAGONAL CYSTINE CRYSTALS ON URINALYSIS, OR 3) POSITIVE FAMILY HISTORY OF CYSTINURIA WITH POSITIVE CYANIDE-NITROPRUSSIDE SCREEN.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	<p>WILSONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST. CYSTINURIA: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST. RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.</p>
<b>Coverage Duration</b>	<p>INITIAL: 12 MONTHS, RENEWAL: LIFETIME.</p>

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: WILSONS DISEASE: 1) KNOWN FAMILY HISTORY OF WILSONS DISEASE OR PHYSICAL EXAMINATION CONSISTENT WITH WILSONS DISEASE, AND 2) REQUESTS FOR FORMULARY VERSION OF PENICILLAMINE CAPSULE REQUIRE A TRIAL OF OR CONTRAINDICATION TO PENICILLAMINE TABLET (DEPEN). CYSTINURIA: REQUESTS FOR FORMULARY VERSION OF PENICILLAMINE CAPSULE REQUIRES A TRIAL OF OR CONTRAINDICATION TO PENICILLAMINE TABLET (DEPEN) AND A FORMULARY VERSION OF TIOPRONIN (THIOLA)/THIOLA EC. RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, 2) TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED, AND 3) REQUESTS FOR FORMULARY VERSION OF PENICILLAMINE CAPSULE REQUIRES A TRIAL OF OR CONTRAINDICATION TO PENICILLAMINE TABLET (DEPEN). RENEWAL: RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, 2) EXPERIENCED OR MAINTAINED IMPROVEMENT IN TENDER JOINT COUNT OR SWOLLEN JOINT COUNT COMPARED TO BASELINE. WILSONS DISEASE, CYSTINURIA: PATIENT CONTINUES TO BENEFIT FROM THE MEDICATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PEXIDARTINIB

---

## Products Affected

- TURALIO ORAL CAPSULE 200 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# PIMAVANSERIN

---

## Products Affected

- NUPLAZID

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	PSYCHOSIS IN PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OR OLDER
<b>Prescriber Restrictions</b>	PSYCHOSIS IN PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, GERIATRICIAN, OR A BEHAVIORAL HEALTH SPECIALIST (E.G., PSYCHIATRIST).
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PSYCHOSIS IN PD: RENEWAL: IMPROVEMENT IN PSYCHOSIS SYMPTOMS FROM BASELINE AND DEMONSTRATES A CONTINUED NEED FOR TREATMENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PIRFENIDONE

## Products Affected

- ESBRIET ORAL CAPSULE
- *pirfenidone oral capsule*
- *pirfenidone oral tablet 267 mg, 534 mg, 801 mg*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	IDIOPATHIC PULMONARY FIBROSIS (IPF): INITIAL: 1) A USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT, AND 2) PREDICTED FORCED VITAL CAPACITY (FVC) OF AT LEAST 50% AT BASELINE.
<b>Age Restrictions</b>	IPF: INITIAL: 18 YEARS OR OLDER.
<b>Prescriber Restrictions</b>	IPF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	IPF: INITIAL: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS, SYSTEMIC SCLEROSIS, RHEUMATOID ARTHRITIS, RADIATION, SARCOIDOSIS, BRONCHIOLITIS OBLITERANS ORGANIZING PNEUMONIA, HUMAN IMMUNODEFICIENCY VIRUS (HIV) INFECTION, VIRAL HEPATITIS, OR CANCER). RENEWAL: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# POMALIDOMIDE

---

## Products Affected

- POMALYST

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PONATINIB

---

## Products Affected

- ICLUSIG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	CML: MUTATIONAL ANALYSIS PRIOR TO INITIATION AND ICLUSIG IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# POSACONAZOLE

---

## Products Affected

- NOXAFIL ORAL SUSPENSION
- *posaconazole*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	OROPHARYNGEAL CANDIDIASIS (OPC): 3 MONTHS. PROPHYLAXIS: 6 MONTHS. TREATMENT: 12 WEEKS.
Other Criteria	POSACONAZOLE SUSPENSION ONLY: 1) OPC: TRIAL OF OR CONTRAINDICATION TO FLUCONAZOLE OR ITRACONAZOLE. 2) PROPHYLAXIS OF INVASIVE ASPERGILLUS AND CANDIDA INFECTION: INABILITY TO SWALLOW TABLETS. POSACONAZOLE TABLETS ONLY: TREATMENT OF INVASIVE ASPERGILLOSIS: NO EXTRA CRITERIA REQUIRED. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# POSACONAZOLE-POWDERMIX

---

## Products Affected

- NOXAFIL ORAL SUSP,DELAYED  
RELEASE FOR RECON

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	6 MONTHS
<b>Other Criteria</b>	PROPHYLAXIS OF INVASIVE ASPERGILLUS AND CANDIDA INFECTION: INABILITY TO SWALLOW TABLETS. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PRALSETINIB

---

## Products Affected

- GAVRETO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# PRAMLINTIDE

---

## Products Affected

- SYMLINPEN 120
- SYMLINPEN 60

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	TYPE I OR TYPE II DIABETES: REQUIRING INSULIN OR CONTINUOUS INSULIN INFUSION (INSULIN PUMP) FOR GLYCEMIC CONTROL
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# PYRIMETHAMINE

## Products Affected

- *pyrimethamine*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	TOXOPLASMOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN INFECTIOUS DISEASE SPECIALIST.
<b>Coverage Duration</b>	TOXOPLASMOSIS: INITIAL: 8 WEEKS, RENEWAL: 6 MOS.
<b>Other Criteria</b>	TOXOPLASMOSIS: RENEWAL: ONE OF THE FOLLOWING: (1) PERSISTENT CLINICAL DISEASE (HEADACHE, NEUROLOGICAL SYMPTOMS, OR FEVER) AND PERSISTENT RADIOGRAPHIC DISEASE (ONE OR MORE MASS LESIONS ON BRAIN IMAGING), OR (2) CD4 COUNT LESS THAN 200 CELLS/MM3 AND CURRENTLY TAKING AN ANTI-RETROVIRAL THERAPY IF HIV POSITIVE.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# QUININE

---

## Products Affected

- *quinine sulfate*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# RAMUCIRUMAB

---

## Products Affected

- CYRAMZA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# REGORAFENIB

---

## Products Affected

- STIVARGA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RELUGOLIX

---

## Products Affected

- ORGOVYX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RESLIZUMAB

---

## Products Affected

- CINQAIR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	ASTHMA: INITIAL: BLOOD EOSINOPHIL LEVEL GREATER THAN OR EQUAL TO 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.
<b>Coverage Duration</b>	ASTHMA: INITIAL 4 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>ASTHMA: INITIAL: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE OF THE FOLLOWING: (A) PATIENT EXPERIENCED AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) PATIENT HAS POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, 3) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: FASENRA, NUCALA, DUPIXENT, AND 4) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA.</p> <p>RENEWAL: 1) NOT CONCURRENTLY RECEIVING XOLAIR, DUPIXENT OR OTHER ANTI-IL5 BIOLOGICS WHEN THESE ARE USED FOR THE TREATMENT OF ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR D) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE. THIS DRUG ALSO REQUIRES</p>
	PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# RIBOCICLIB

## Products Affected

- KISQALI FEMARA CO-PACK ORAL TABLET 200 MG/DAY(200 MG X 1)-2.5 MG, 400 MG/DAY(200 MG X 2)-2.5 MG, 600 MG/DAY(200 MG X 3)-2.5 MG
- KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ADVANCED OR METASTATIC BREAST CANCER: TRIAL OF OR CONTRAINDICATION TO VERZENIO OR IBRANCE WHERE INDICATIONS ALIGN.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# RIFAXIMIN

---

## Products Affected

- XIFAXAN ORAL TABLET 200 MG,  
550 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	TRAVELERS DIARRHEA/HE: 12 MOS. IBS-D: 8 WKS.
<b>Other Criteria</b>	RIFAXIMIN 550 MG TABLETS: HE: TRIAL OF OR CONTRAINDICATION TO LACTULOSE OR CONCURRENT LACTULOSE THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RIMEGEPANT

## Products Affected

- NURTEC ODT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	<p>INITIAL: ACUTE MIGRAINE TREATMENT: TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN). EPISODIC MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREVENTIVE MIGRAINE TREATMENTS: DIVALPROEX SODIUM, TOPIRAMATE, PROPRANOLOL, TIMOLOL. RENEWAL: ACUTE MIGRAINE TREATMENT: 1) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR 2) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS. EPISODIC MIGRAINE PREVENTION: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION, AND 2) REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY.</p>
Indications	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RIOCIQUAT

---

## Products Affected

- ADEMPAS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PULMONARY ARTERIAL HYPERTENSION (PAH): 1) CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER, AND 2) NYHA-WHO FUNCTIONAL CLASS (FC) II-IV SYMPTOMS. PERSISTENT/RECURRENT CHRONIC THROMBOEMBOLIC PULMONARY HYPERTENSION (CTEPH) WHO GROUP 4: NYHA-WHO FUNCTIONAL CLASS II-IV SYMPTOMS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PAH, CTEPH: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL AND RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: PAH: NOT CONCURRENTLY TAKING NITRATES, NITRIC OXIDE DONORS, PHOSPHODIESTERASE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS. CTEPH: 1) NOT CONCURRENTLY TAKING NITRATES, NITRIC OXIDE DONORS, OR ANY PDE INHIBITORS, AND 2) NOT A CANDIDATE FOR SURGERY OR HAS INOPERABLE CTEPH OR HAS PERSISTENT OR RECURRENT DISEASE AFTER SURGICAL TREATMENT. RENEWAL: PAH, CTEPH: 1) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE OR A STABLE 6-MINUTE WALK DISTANCE WITH A STABLE/IMPROVED WHO FUNCTIONAL CLASS, AND 2) NOT CONCURRENTLY TAKING NITRATES, NITRIC OXIDE DONORS, PHOSPHODIESTERASE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RIPRETINIB

---

## Products Affected

- QINLOCK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# RISANKIZUMAB-RZAA

## Products Affected

- SKYRIZI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PLAQUE PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE OR GENITAL AREA.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). CD: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPYRINE, METHOTREXATE, OR MESALAMINE. RENEWAL: PSO, PSA: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RISDIPLAM

## Products Affected

- EVRYSDI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	SPINAL MUSCULAR ATROPHY (SMA): INITIAL: GENE MUTATION ANALYSIS INDICATING MUTATIONS OR DELETIONS OF BOTH ALLELES OF THE SURVIVAL MOTOR NEURON 1 (SMN1) GENE. FOR PRESYMPTOMATIC PATIENTS: UP TO THREE COPIES OF SURVIVAL MOTOR NEURON 2 (SMN2) BASED ON NEWBORN SCREENING.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	SMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROMUSCULAR SPECIALIST OR SMA SPECIALIST AT A SMA SPECIALTY CENTER.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS
<b>Other Criteria</b>	SMA: INITIAL: FOR SYMPTOMATIC PATIENTS: (1) BASELINE MOTOR FUNCTION ASSESSMENT BY A NEUROMUSCULAR SPECIALIST OR SMA SPECIALIST, AND (2) IF PATIENT RECEIVED GENE THERAPY, THE PATIENT HAD LESS THAN EXPECTED CLINICAL BENEFIT. RENEWAL: IMPROVED, MAINTAINED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN: (1) MOTOR FUNCTION ASSESSMENTS COMPARED TO BASELINE, OR (2) OTHER MUSCLE FUNCTION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RITUXIMAB AND HYALURONIDASE HUMAN-SQ

## Products Affected

- RITUXAN HYCELA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	PATIENT HAS RECEIVED OR WILL RECEIVE AT LEAST ONE FULL DOSE OF A RITUXIMAB PRODUCT BY INTRAVENOUS INFUSION PRIOR TO INITIATION OF RITUXIMAB AND HYALURONIDASE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# RITUXIMAB-ABBS

## Products Affected

- TRUXIMA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NHL, CLL: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.
<b>Coverage Duration</b>	RA: INITIAL: 6 MO, RENEWAL: 12 MO. NHL, GPA, MPA: 12 MO. CLL: 6 MO.
<b>Other Criteria</b>	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RITUXIMAB-ARRX

## Products Affected

- RIABNI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NHL, CLL: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.
<b>Coverage Duration</b>	RA: INITIAL: 6 MO, RENEWAL: 12 MO. NHL, WG, MPA: 12 MO. CLL: 6 MO.
<b>Other Criteria</b>	RA: INITIAL: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RITUXIMAB-PVVR

## Products Affected

- RUXIENCE

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NHL, CLL: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.
<b>Coverage Duration</b>	NHL, GPA, MPA: 12 MONTHS. CLL: 6 MONTHS. RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO PREFERRED AGENTS: HUMIRA, ENBREL, RINVOQ, XELJANZ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ROMIPLOSTIM

---

## Products Affected

- NPLATE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	IMMUNE THROMBOCYTOPENIA (ITP): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.
<b>Coverage Duration</b>	ITP: INITIAL: 4 MO, RENEWAL: 12 MO. HEMATOPOIETIC SYNDROME OF ACUTE RADIATION SYNDROME: 12 MO.
<b>Other Criteria</b>	ITP: INITIAL: TRIAL OF OR CONTRAINDICATION TO CORTICOSTEROIDS, IMMUNOGLOBULINS, OR AN INSUFFICIENT RESPONSE TO SPLENECTOMY. RENEWAL: CLINICAL RESPONSE TO THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# ROMOSOZUMAB

## Products Affected

- EVENITY

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	<p>POSTMENOPAUSAL OSTEOPOROSIS: 1) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF EVENITY TREATMENT, 2) ONE OF THE FOLLOWING: (A) HIGH RISK FOR FRACTURES DEFINED AS ONE OF THE FOLLOWING: (I) HISTORY OF OSTEOPOROTIC (I.E., FRAGILITY, LOW TRAUMA) FRACTURE(S), (II) 2 OR MORE RISK FACTORS FOR FRACTURE (E.G., HISTORY OF MULTIPLE RECENT LOW TRAUMA FRACTURES, BMD T-SCORE LESS THAN OR EQUAL TO -2.5, CORTICOSTEROID USE, USE OF GNRH ANALOGS SUCH AS NAFARELIN, ETC.), (III) NO PRIOR TREATMENT FOR OSTEOPOROSIS AND FRAX SCORE OF AT LEAST 20% FOR ANY MAJOR FRACTURE OR OF AT LEAST 3% FOR HIP FRACTURE, (B) UNABLE TO USE ORAL THERAPY (I.E., UPPER GASTROINTESTINAL PROBLEMS - UNABLE TO TOLERATE ORAL MEDICATION, LOWER GASTROINTESTINAL PROBLEMS - UNABLE TO ABSORB ORAL MEDICATIONS, TROUBLE REMEMBERING TO TAKE ORAL MEDICATIONS OR COORDINATING AN ORAL BISPHOSPHONATE WITH OTHER ORAL MEDICATIONS OR THEIR DAILY ROUTINE), OR (C) TRIAL OF, INTOLERANCE TO, OR A CONTRAINDICATION TO A BISPHOSPHONATE.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# ROPEGINTERFERON ALFA-2B-NJFT

---

## Products Affected

- BESREMI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RUCAPARIB

---

## Products Affected

- RUBRACA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: ONE OF THE FOLLOWING: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# RUXOLITINIB

---

## Products Affected

- JAKAFI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	MYELOFIBROSIS: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. POLYCYTHEMIA VERA, GVHD: 12 MONTHS.
<b>Other Criteria</b>	MYELOFIBROSIS: RENEWAL: PATIENT CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SAFINAMIDE

---

## Products Affected

- XADAGO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SAPROPTERIN

---

## Products Affected

- *javygtor oral tablet, soluble*
- *sapropterin oral tablet, soluble*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	INITIAL: 1 MONTH, RENEWAL 12 MONTHS.
Other Criteria	HYPERPHENYLALANINEMIA (HPA): INITIAL: NOT CONCURRENTLY USING PALYNZIQ. RENEWAL: 1) CONTINUES TO BENEFIT FROM TREATMENT, AND 2) NOT CONCURRENTLY USING PALYNZIQ.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SARILUMAB

## Products Affected

- KEVZARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	RHEUMATOID ARTHRITIS (RA): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
Other Criteria	RA: INITIAL: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# SATRALIZUMAB-MWGE

## Products Affected

- ENSPRYNG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	NMOSD: INITIAL: PRESCRIBED BY AN OPHTHALMOLOGIST OR PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	NMOSD: INITIAL: 1) ONE OF THE FOLLOWING CORE CLINICAL CHARACTERISTIC: (A) OPTIC NEURITIS, (B) ACUTE MYELITIS, (C) AREA POSTREMA SYNDROME, (D) ACUTE BRAINSTEM SYNDROME, (E) SYMPTOMATIC NARCOLEPSY OR ACUTE DIENCEPHALIC CLINICAL SYNDROME WITH NMOSD-TYPICAL DIENCEPHALIC MRI LESIONS, OR (F) SYMPTOMATIC CEREBRAL SYNDROME WITH NMOSD-TYPICAL BRAIN LESIONS, AND 2) NO CONCURRENT USE WITH RITUXIMAB, INEBILIZUMAB, OR ECULIZUMAB. RENEWAL: 1) REDUCTION IN RELAPSE FREQUENCY FROM BASELINE, AND 2) NO CONCURRENT USE WITH USE RITUXIMAB, INEBILIZUMAB, OR ECULIZUMAB.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SEBELIPASE ALFA

## Products Affected

- KANUMA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	<p>LYSOSOMAL ACID LIPASE (LAL) DEFICIENCY: INITIAL: DIAGNOSIS CONFIRMED BY 1) PRESENCE OF CLINICAL FEATURES (E.G., HEPATOMEGALY, ELEVATED SERUM TRANSAMINASES, DYSLIPIDEMIA, SPLENOMEGALY), AND 2) ONE OF THE FOLLOWING: (A) BLOOD TEST INDICATING LOW OR ABSENT LEVELS OF LAL ENZYME ACTIVITY, (B) DRIED BLOOD SPOT TEST INDICATING LOW OR ABSENT LAL ENZYME ACTIVITY, OR (C) GENETIC TEST INDICATING THE BI-ALLELIC PRESENCE OF ALTERED LIPA GENE(S).</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	<p>LAL DEFICIENCY: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST, HEPATOLOGIST, GASTROENTEROLOGIST, MEDICAL GENETICIST, LIPIDOLOGIST, OR METABOLIC SPECIALIST.</p>
<b>Coverage Duration</b>	<p>LAL DEFICIENCY: INITIAL: 6 OR 12 MONTHS, SEE OTHER CRITERIA. RENEWAL: 12 MONTHS</p>

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	<p>LAL DEFICIENCY: INITIAL: RAPIDLY PROGRESSIVE LAL DEFICIENCY PRESENTING WITHIN THE FIRST 6 MONTHS OF LIFE: 12 MONTHS. LAL DEFICIENCY PRESENTING AFTER THE FIRST 6 MONTHS OF LIFE AND NOT CONSIDERED RAPIDLY PROGRESSIVE: 6 MONTHS. RENEWAL: LAL DEFICIENCY PRESENTING AFTER THE FIRST 6 MONTHS OF LIFE AND NOT CONSIDERED RAPIDLY PROGRESSIVE: IMPROVEMENT IN ONE OF THE FOLLOWING CLINICAL PARAMETERS DURING THE PAST 6 MONTHS: (1) A RELATIVE REDUCTION FROM BASELINE IN LDL-C, NON-HDL-C, OR TRIGLYCERIDES, (2) NORMALIZATION OF ASPARTATE AMINOTRANSFERASE (AST) BASED ON AGE- AND GENDER-SPECIFIC NORMAL RANGES, OR (3) DECREASE IN LIVER FAT CONTENT COMPARED TO BASELINE ASSESSED BY ABDOMINAL IMAGING.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SECUKINUMAB

## Products Affected

- COSENTYX (2 SYRINGES)
- COSENTYX PEN (2 PENS)
- COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR A DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA, ENTHESITIS-RELATED ARTHRITIS (ERA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS

PA Criteria	Criteria Details
<b>Other Criteria</b>	INITIAL: PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTI-RHEUMATIC DRUG). AS, NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID (NON-STEROIDAL ANTI-INFLAMMATORY DRUG). ERA: TRIAL OF OR CONTRAINDICATION TO ONE NSAID, SULFASALAZINE, OR METHOTREXATE. RENEWAL: PSO, PSA, AS, NR-AXSPA, ERA: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SELEXIPAG

## Products Affected

- UPTRAVI INTRAVENOUS
- UPTRAVI ORAL TABLET 1,000 MCG, 1,200 MCG, 1,400 MCG, 1,600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG
- UPTRAVI ORAL TABLETS,DOSE PACK

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: 1) CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER, AND 2) NYHA-WHO FUNCTIONAL CLASS (FC) II-IV SYMPTOMS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>PAH: INITIAL: WHO FC II-III SYMPTOMS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING AGENTS, EACH FROM A DIFFERENT DRUG CLASS: 1) FORMULARY VERSION OF AN ORAL ENDOTHELIAL RECEPTOR ANTAGONIST, 2) FORMULARY VERSION OF AN ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR, OR 3) FORMULARY VERSION OF AN ORAL CGMP STIMULATOR. WHO FC III SYMPTOMS AND EVIDENCE OF RAPID PROGRESSION OR POOR PROGNOSIS, WHO FC IV SYMPTOMS: NO STEP. RENEWAL: 1) IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST, OR 2) REMAINED STABLE FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST AND WHO FC HAS IMPROVED OR REMAINED STABLE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SELINEXOR

---

## Products Affected

- XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40 MG/WEEK (40 MG X 1), 40MG TWICE WEEK (40 MG X 2), 60 MG/WEEK (60 MG X 1), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (40 MG X 2), 80MG TWICE WEEK (160 MG/WEEK)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No



# SELPERCATINIB

---

## Products Affected

- RETEVMO ORAL CAPSULE 40 MG,  
80 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SELUMETINIB

---

## Products Affected

- KOSELUGO ORAL CAPSULE 10 MG,  
25 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SIPONIMOD

## Products Affected

- MAYZENT ORAL TABLET 0.25 MG, 1 MG, 2 MG
- MAYZENT STARTER(FOR 1MG MAINT)
- MAYZENT STARTER(FOR 2MG MAINT)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	MULTIPLE SCLEROSIS: RENEWAL: 1) DEMONSTRATION OF CLINICAL BENEFIT COMPARED TO PRE-TREATMENT BASELINE AND 2) DOES NOT HAVE LYMPHOPENIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SIROLIMUS PROTEIN-BOUND

---

## Products Affected

- FYARRO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SODIUM OXYBATE

## Products Affected

- *sodium oxybate*
- XYREM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: ALL INDICATIONS: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR SPECIALIST IN SLEEP MEDICINE.
Coverage Duration	INITIAL 6 MONTHS. RENEWAL: 12 MONTHS
Other Criteria	INITIAL: EXCESSIVE DAYTIME SLEEPINESS (EDS) IN NARCOLEPSY: 1) NOT CURRENTLY TAKING A SEDATIVE HYPNOTIC AGENT, 2) FOR PATIENTS 18 YEARS OR OLDER: TRIAL OF OR CONTRAINDICATION TO THE FORMULARY VERSION OF MODAFINIL, ARMODAFINIL, PITOLISANT OR SOLRIAMFETOL AND ONE OTHER GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY, AND 3) FOR PATIENTS 7 TO 17 YEARS OF AGE: TRIAL OF OR CONTRAINDICATION TO ONE OTHER GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY. CATAPLEXY IN NARCOLEPSY: NOT CURRENTLY TAKING A SEDATIVE HYPNOTIC AGENT. RENEWAL (ALL INDICATIONS): 1) SUSTAINED IMPROVEMENT OF SYMPTOMS COMPARED TO BASELINE, AND 2) NOT CURRENTLY TAKING A SEDATIVE HYPNOTIC AGENT.
Indications	All FDA-approved Indications.
Off Label Uses	

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# SOFOSBUVIR/VELPATASVIR

## Products Affected

- EPCLUSA ORAL PELLETS IN PACKET 150-37.5 MG, 200-50 MG
- EPCLUSA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	HCV RNA LEVEL WITHIN PAST 6 MONTHS.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
Other Criteria	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, HIV REGIMEN THAT CONTAINS EFAVIRENZ, ROSUVASTATIN AT DOSES ABOVE 10MG, TIPRANA VIR/RITONAVIR, TOPOTECAN, SOVALDI (AS A SINGLE AGENT), HARVONI, ZEPATIER, MAVYRET, OR VOSEVI, AND 3) PATIENTS WITH DECOMPENSATED CIRRHOSIS REQUIRE CONCURRENT RIBAVIRIN UNLESS RIBAVIRIN INELIGIBLE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SOFOSBUVIR/VELPATASVIR/VOXILAPREVIR

## Products Affected

- VOSEVI

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	HCV RNA LEVEL WITHIN PAST 6 MONTHS
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.
<b>Other Criteria</b>	1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, CYCLOSPORINE, PITAVASTATIN, PRAVASTATIN (DOSES ABOVE 40MG), ROSUVASTATIN, METHOTREXATE, MITOXANTRONE, IMATINIB, IRINOTECAN, LAPATINIB, SULFASALAZINE, TOPOTECAN, OR HIV REGIMEN THAT CONTAINS EFAVIRENZ, ATAZANAVIR, LOPINAVIR, TIPRANAVIR/RITONAVIR, SOVALDI (AS A SINGLE AGENT), EPCLUSA, HARVONI, ZEPATIER, OR MAVYRET, AND 3) DOES NOT HAVE MODERATE OR SEVERE HEPATIC IMPAIRMENT (CHILD-PUGH B OR C).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Part B Prerequisite</b>	No

# SOLRIAMFETOL

---

## Products Affected

- SUNOSI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: EXCESSIVE DAYTIME SLEEPINESS (EDS) IN NARCOLEPSY: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR SPECIALIST IN SLEEP MEDICINE.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: EDS IN NARCOLEPSY: TRIED THE FORMULARY VERSION OF MODAFINIL OR ARMODAFINIL, AND ONE OTHER GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY. EDS IN OBSTRUCTIVE SLEEP APNEA (OSA): TRIED THE FORMULARY VERSION OF MODAFINIL OR ARMODAFINIL. RENEWAL: EDS IN NARCOLEPSY OR OSA: SUSTAINED IMPROVEMENT OF SYMPTOMS COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SOMATROPIN - NORDITROPIN

---

## Products Affected

- NORDITROPIN FLEXPPO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES.
<b>Required Medical Information</b>	INITIAL: PEDIATRIC GROWTH HORMONE DEFICIENCY (GHD), IDIOPATHIC SHORT STATURE (ISS), SMALL FOR GESTATIONAL AGE (SGA), TURNER SYNDROME (TS), NOONAN SYNDROME: HEIGHT AT LEAST 2 STANDARD DEVIATIONS (SD) BELOW THE MEAN HEIGHT FOR NORMAL CHILDREN OF THE SAME AGE AND GENDER. PRADER WILLI SYNDROME (PWS): CONFIRMED GENETIC DIAGNOSIS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL/RENEWAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: ADULT GHD: GROWTH HORMONE DEFICIENCY ALONE OR ASSOCIATED WITH MULTIPLE HORMONE DEFICIENCIES (HYPOPITUITARISM), AS A RESULT OF PITUITARY DISEASES, HYPOTHALAMIC DISEASE, SURGERY, RADIATION THERAPY, TRAUMA, OR CONTINUATION OF THERAPY FROM CHILDHOOD ONSET GROWTH HORMONE DEFICIENCY. PEDIATRIC GHD, ISS, SGA, TS, NOONAN SYNDROME: OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND. RENEWAL: PEDIATRIC GHD: 1) IMPROVEMENT WHILE ON THERAPY (I.E., INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND OR PATIENT HAS NOT COMPLETED PREPUBERTALGROWTH. ISS, SGA, TS, NOONAN SYNDROME: 1) IMPROVEMENT WHILE ON THERAPY (I.E., INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND. PWS: IMPROVEMENT IN BODY COMPOSITION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SOMATROPIN - SEROSTIM

## Products Affected

- SEROSTIM SUBCUTANEOUS  
RECON SOLN 4 MG, 5 MG, 6 MG

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES
<b>Required Medical Information</b>	INITIAL: HIV/WASTING: MEETS ONE OF THE FOLLOWING CRITERIA FOR WEIGHT LOSS: 1) 10% UNINTENTIONAL WEIGHT LOSS OVER 12 MONTHS, 2) 7.5% UNINTENTIONAL WEIGHT LOSS OVER 6 MONTHS, 3) 5% BODY CELL MASS (BCM) LOSS WITHIN 6 MONTHS, 4) BODY CELL MASS (BCM) LESS THAN 35% (MEN) OF TOTAL BODY WEIGHT AND BODY MASS INDEX (BMI) LESS THAN 27 KG PER METER SQUARED, 5) BCM LESS THAN 23% (WOMEN) OF TOTAL BODY WEIGHT AND A BODY MASS INDEX (BMI) LESS THAN 27 KG PER METER SQUARED, OR 6) BMI LESS THAN 18.5 KG PER METER SQUARED.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	HIV/WASTING: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, NUTRITIONAL SUPPORT SPECIALIST, OR INFECTIOUS DISEASE SPECIALIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 3 MONTHS.
<b>Other Criteria</b>	HIV/WASTING: INITIAL: 1) CURRENTLY ON HIV ANTIRETROVIRAL THERAPY, AND 2) INADEQUATE RESPONSE TO ONE PREVIOUS THERAPY (E.G., MEGACE, APPETITE STIMULANTS, ANABOLIC STEROIDS). RENEWAL: 1) CURRENTLY ON HIV ANTIRETROVIRAL THERAPY, AND 2) CLINICAL BENEFIT IN MUSCLE MASS AND WEIGHT.
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SONIDEGIB

---

## Products Affected

- ODOMZO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	LOCALLY ADVANCED BASAL CELL CARCINOMA (BCC); BASELINE SERUM CREATINE KINASE (CK) AND SERUM CREATININE LEVELS
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# SORAFENIB

---

## Products Affected

- *sorafenib*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# SOTORASIB

---

## Products Affected

- LUMAKRAS

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# STIRIPENTOL

---

## Products Affected

- DIACOMIT ORAL CAPSULE 250 MG, 500 MG
- DIACOMIT ORAL POWDER IN PACKET 250 MG, 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	DRAVET SYNDROME: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
Coverage Duration	INITIAL/RENEWAL: 12 MONTHS.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SUNITINIB

---

## Products Affected

- *sunitinib*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	GASTROINTESTINAL STROMAL TUMORS (GIST): TRIAL OF OR CONTRAINDICATION TO IMATINIB (GLEEVEC).
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TADALAFIL

---

## Products Affected

- *tadalafil oral tablet 2.5 mg, 5 mg*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	ERECTILE DYSFUNCTION WITHOUT DIAGNOSIS OF BENIGN PROSTATIC HYPERPLASIA (BPH).
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	BPH: 1) TRIAL OF ONE ALPHA BLOCKER (SUCH AS DOXAZOSIN, TERAZOSIN, TAMSULOSIN, ALFUZOSIN), AND 2) TRIAL OF ONE 5-ALPHA-REDUCTASE INHIBITOR (SUCH AS FINASTERIDE, DUTASTERIDE). APPLIES TO 2.5MG AND 5MG STRENGTHS ONLY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TALAZOPARIB

---

## Products Affected

- TALZENNA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	PATIENT HAS BEEN TREATED WITH CHEMOTHERAPY IN THE NEOADJUVANT, ADJUVANT, OR METASTATIC SETTING. PATIENTS WITH HORMONE RECEPTOR (HR)-POSITIVE BREAST CANCER MUST HAVE ADDITIONAL PRIOR TREATMENT WITH ENDOCRINE THERAPY OR BE CONSIDERED INAPPROPRIATE FOR ENDOCRINE THERAPY.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TALIMOGENE

## Products Affected

- IMLYGIC INJECTION SUSPENSION  
10EXP6 (1 MILLION) PFU/ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	UNRESECTABLE MELANOMA: 1) IMLYGIC TO BE INJECTED INTO CUTANEOUS, SUBCUTANEOUS, AND/OR NODAL LESIONS THAT ARE VISIBLE, PALPABLE, OR DETECTABLE BY ULTRASOUND GUIDANCE, 2) NOT CURRENTLY RECEIVING IMMUNOSUPPRESSIVE THERAPY, 3) NO HISTORY OF PRIMARY OR ACQUIRED IMMUNODEFICIENT STATES, LEUKEMIA, LYMPHOMA, OR AIDS, AND 4) NO CONCURRENT USE WITH PEMBROLIZUMAB, NIVOLUMAB, IPILIMUMAB, DABRAFENIB, TRAMETINIB, VEMURAFENIB, INTERLEUKIN-2, INTERFERON, DACARBAZINE, TEMOZOLOMIDE, PACLITAXEL, CARBOPLATIN, IMATINIB, MELPHALAN, IMIQUIMOD, OR RADIATION THERAPY.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TASIMELTEON

---

## Products Affected

- HETLIOZ
- HETLIOZ LQ
- *tasimelteon*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	LIFETIME
<b>Other Criteria</b>	NON-24 HOUR SLEEP-WAKE DISORDER: PATIENT IS LIGHT-INSENSITIVE OR HAS TOTAL BLINDNESS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TAZEMETOSTAT

---

## Products Affected

- TAZVERIK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# TBO-FILGRASTIM

---

## Products Affected

- GRANIX

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	NON-MYELOID MALIGNANCIES: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	NON-MYELOID MALIGNANCIES: TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: NIVESTYM.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TEBENTAFUSP-TEBN

---

## Products Affected

- KIMMTRAK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TECLISTAMAB-CQYV

---

## Products Affected

- TECVAYLI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TEDUGLUTIDE

---

## Products Affected

- GATTEX 30-VIAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	SHORT BOWEL SYNDROME (SBS): INITIAL/RENEWAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS
Other Criteria	SBS: INITIAL: PATIENT IS DEPENDENT ON INTRAVENOUS PARENTERAL NUTRITION DEFINED AS REQUIRING PARENTERAL NUTRITION AT LEAST THREE TIMES PER WEEK. RENEWAL: ACHIEVED OR MAINTAINED A DECREASED NEED FOR PARENTERAL SUPPORT COMPARED TO BASELINE.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TELOTRISTAT

---

## Products Affected

- XERMELO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CARCINOID SYNDROME DIARRHEA: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST OR GASTROENTEROLOGIST
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TEPOTINIB

---

## Products Affected

- TEPMETKO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TEPROTUMUMAB-TRBW

---

## Products Affected

- TEPEZZA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TERIFLUNOMIDE

---

## Products Affected

- AUBAGIO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# TERIPARATIDE

---

## Products Affected

- FORTEO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>OSTEOPOROSIS: ONE OF THE FOLLOWING: (1) HIGH RISK FOR FRACTURES DEFINED AS ONE OF THE FOLLOWING: A) HISTORY OF OSTEOPOROTIC (I.E., FRAGILITY, LOW TRAUMA) FRACTURE(S), B) 2 OR MORE RISK FACTORS FOR FRACTURE (E.G., HISTORY OF MULTIPLE RECENT LOW TRAUMA FRACTURES, BMD T-SCORE LESS THAN OR EQUAL TO -2.5, CORTICOSTEROID USE, OR USE OF GNRH ANALOGS SUCH AS NAFARELIN, ETC.), C) NO PRIOR TREATMENT FOR OSTEOPOROSIS AND FRAX SCORE OF AT LEAST 20% FOR ANY MAJOR FRACTURE OR OF AT LEAST 3% FOR HIP FRACTURE, (2) UNABLE TO USE ORAL THERAPY (I.E., UPPER GASTROINTESTINAL PROBLEMS UNABLE TO TOLERATE ORAL MEDICATION, LOWER GASTROINTESTINAL PROBLEMS UNABLE TO ABSORB ORAL MEDICATIONS, TROUBLE REMEMBERING TO TAKE ORAL MEDICATIONS OR COORDINATING AN ORAL BISPHOSPHONATE WITH OTHER ORAL MEDICATIONS OR THEIR DAILY ROUTINE), (3) TRIAL OF, INTOLERANCE TO, OR A CONTRAINDICATION TO ONE BISPHOSPHONATE. PATIENT HAS NOT RECEIVED A TOTAL OF 24 MONTHS CUMULATIVE TREATMENT UNLESS PATIENT REMAINS AT OR HAS RETURNED TO HAVING A HIGH RISK FOR FRACTURE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TESAMORELIN

---

## Products Affected

- EGRIFTA SV

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	3 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TESTOSTERONE

## Products Affected

- *testosterone cypionate*
- *testosterone enanthate*
- *testosterone transdermal gel in metered-dose pump 12.5 mg/1.25 gram (1%), 20.25 mg/1.25 gram (1.62%)*
- *testosterone transdermal gel in packet 1% (25 mg/2.5gram), 1% (50 mg/5 gram)*
- *testosterone transdermal solution in metered pump w/lapp*
- XYOSTED

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: MALE HYPOGONADISM: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	MALE HYPOGONADISM: INITIAL/RENEWAL:12 MO. ALL OTHER INDICATIONS: LIFETIME OF MEMBERSHIP IN PLAN.
<b>Other Criteria</b>	RENEWAL: MALE HYPOGONADISM: IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TETRABENAZINE

---

## Products Affected

- *tetrabenazine*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	HUNTINGTONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST
Coverage Duration	12 MONTHS
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TEZACAFTOR/IVACAFTOR

## Products Affected

- SYMDEKO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	CYSTIC FIBROSIS (CF); INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: LIFETIME
<b>Other Criteria</b>	CF: RENEWAL: MAINTAINED, IMPROVED, OR DEMONSTRATED LESS THAN EXPECTED DECLINE IN FEV1 OR BODY MASS INDEX (BMI), OR REDUCTION IN NUMBER OF PULMONARY EXACERBATIONS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# THALIDOMIDE

---

## Products Affected

- THALOMID

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TILDRAKIZUMAB-ASMN

---

## Products Affected

- ILUMYA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PLAQUE PSORIASIS (PSO): INITIAL: PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PSO: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	PSO: INITIAL: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, COSENTYX, STELARA, ENBREL, SKYRIZI, TREMFYA. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# TISOTUMAB VEDOTIN-TFTV

---

## Products Affected

- TIVDAK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TIVOZANIB

---

## Products Affected

- FOTIVDA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TOCILIZUMAB IV

## Products Affected

- ACTEMRA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST.
<b>Coverage Duration</b>	INITIAL: RA, PJIA, SJIA, GCA: 6 MONTHS. CRS: 1 MONTH. RENEWAL: RA, PJIA, SJIA, GCA: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. SJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). RENEWAL: RA, PJIA, SJIA: CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TOCILIZUMAB SQ

---

## Products Affected

- ACTEMRA
- ACTEMRA ACTPEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST. SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE (SSC-ILD): PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR RHEUMATOLOGIST.
Coverage Duration	INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: ONE OF THE FOLLOWING: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ, RINVOQ, OR 2) TRIAL OF A TNF INHIBITOR AND PHYSICIAN HAS INDICATED THE PATIENT CANNOT USE A JAK INHIBITOR DUE TO THE BLACK BOX WARNING FOR INCREASED RISK OF MORTALITY, MALIGNANCIES, AND SERIOUS CARDIOVASCULAR EVENTS. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: HUMIRA, ENBREL, XELJANZ IR. SJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). SSC-ILD: DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS). RENEWAL: RA, PJIA, SJIA: CONTINUES TO BENEFIT FROM THE MEDICATION. SSC-ILD: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TOFACITINIB

## Products Affected

- XELJANZ
- XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), POLYARTICULAR COURSE JUVENILE IDIOPATHIC ARTHRITIS (PCJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PSA, PCJIA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. AS: TRIAL OF OR CONTRAINDICATION TO AN NSAID. UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPYRINE, METHOTREXATE, OR MESALAMINE. RENEWAL: RA, PSA, AS, PCJIA: CONTINUES TO BENEFIT FROM THE MEDICATION.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# TOLVAPTAN

## Products Affected

- JYNARQUE ORAL TABLET
- JYNARQUE ORAL TABLETS, SEQUENTIAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	AUTOSOMAL DOMINANT POLYCYSTIC KIDNEY DISEASE (ADPKD): INITIAL: 1) CONFIRMED POLYCYSTIC KIDNEY DISEASE VIA CT, MRI, OR ULTRASOUND, AND 2) GENETIC TESTING FOR CAUSATIVE MUTATIONS OR FAMILY HISTORY OF CONFIRMED POLYCYSTIC KIDNEY DISEASE IN ONE OR BOTH PARENTS.
Age Restrictions	
Prescriber Restrictions	ADPKD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	ADPKD: INITIAL: DOES NOT HAVE ESRD (I.E., RECEIVING DIALYSIS OR HAS UNDERGONE RENAL TRANSPLANT). RENEWAL: HAS NOT PROGRESSED TO ESRD/DIALYSIS OR TRANSPLANT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TOPICAL TRETINOIN

---

## Products Affected

- ALTRENO
- *tretinoin*

PA Criteria	Criteria Details
Exclusion Criteria	COSMETIC INDICATIONS SUCH AS WRINKLES, PHOTOAGING, MELASMA.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	12 MONTHS
Other Criteria	ACNE VULGARIS: BRAND TOPICAL TRETINOIN REQUIRES TRIAL OF OR CONTRAINDICATION TO A GENERIC TOPICAL TRETINOIN PRODUCT.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TRAMETINIB

---

## Products Affected

- MEKINIST ORAL TABLET 0.5 MG, 2 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRASTUZUMAB HYALURONIDASE

---

## Products Affected

- HERCEPTIN HYLECTA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	ADJUVANT BREAST CANCER, METASTATIC BREAST CANCER: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: HERZUMA, OGIVRI, ONTRUZANT, TRAZIMERA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRASTUZUMAB-ANNS

---

## Products Affected

- KANJINTI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	ADJUVANT BREAST CANCER, METASTATIC BREAST CANCER: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: HERZUMA, OGIVRI, ONTRUZANT, TRAZIMERA. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRASTUZUMAB-DKST

---

## Products Affected

- OGIVRI

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRASTUZUMAB-DTTB

---

## Products Affected

- ONTRUZANT

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRASTUZUMAB-PKRB

---

## Products Affected

- HERZUMA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# TRASTUZUMAB-QYYP

---

## Products Affected

- TRAZIMERA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TREMELIMUMAB-ACTL

---

## Products Affected

- IMJUDO

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	UHCC: 30 DAYS. METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC): 5 MONTHS.
<b>Other Criteria</b>	UNRESECTABLE HEPATOCELLULAR CARCINOMA (UHCC): HAS NOT RECEIVED PRIOR TREATMENT WITH IMJUDO.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TREPROSTINIL INHALED

## Products Affected

- TYVASO

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	<p>INITIAL: PULMONARY ARTERIAL HYPERTENSION (PAH): (1) CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER, AND (2) NYHA-WHO FUNCTIONAL CLASS (FC) III-IV SYMPTOMS. PULMONARY HYPERTENSION-INTERSTITIAL LUNG DISEASE (PH-ILD): CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION: (1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, (2) PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER, AND (3) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS.</p>
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	<p>INITIAL: PAH, PH-ILD: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.</p>
<b>Coverage Duration</b>	<p>INITIAL: PAH: 12 MONTHS, PH-ILD: 6 MONTHS. RENEWAL: PAH, PH-ILD: 12 MONTHS.</p>

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: PAH: WHO FC III SYMPTOMS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING AGENTS FROM DIFFERENT DRUG CLASSES: 1) FORMULARY VERSION OF AN ORAL ENDOTHELIAL RECEPTOR ANTAGONIST, 2) FORMULARY VERSION OF AN ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR, OR 3) FORMULARY VERSION OF AN ORAL CGMP STIMULATOR. WHO FC III SYMPTOMS AND EVIDENCE OF RAPID PROGRESSION OR POOR PROGNOSIS, WHO FC IV SYMPTOMS: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF AN IV/SQ PROSTACYCLIN. RENEWAL: PAH: IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST OR REMAINED STABLE FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST AND WHO FC HAS IMPROVED OR REMAINED STABLE. PH-ILD: IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST OR A STABLE 6-MINUTE WALK DISTANCE TEST. THIS DRUG MAY BE COVERED UNDER MEDICARE PART B OR D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TREPROSTINIL INJECTABLE

## Products Affected

- *treprostinil sodium*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: (1) CONFIRMATORY DIAGNOSIS BASED ON RIGHT HEART CATHETERIZATION: A) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, B) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) 15 MMHG OR LESS, AND C) PULMONARY VASCULAR RESISTANCE (PVR) 3 WOOD UNITS OR GREATER, AND (2) NYHA-WHO FUNCTIONAL CLASS (FC) II-IV SYMPTOMS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.
<b>Coverage Duration</b>	PAH: INITIAL AND RENEWAL: 12 MONTHS
<b>Other Criteria</b>	PAH: INITIAL: CONTINUATION FROM HOSPITAL DISCHARGE FOR WHO FC II-IV: NO STEP. NEW STARTS FOR WHO FC III-IV: NO STEP. NEW STARTS FOR WHO FC II: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING AGENTS FROM DIFFERENT DRUG CLASSES: 1) FORMULARY VERSION OF AN ORAL ENDOTHELIAL RECEPTOR ANTAGONIST, 2) FORMULARY VERSION OF AN ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR, OR 3) FORMULARY VERSION OF AN ORAL CGMP STIMULATOR. RENEWAL: IMPROVEMENT FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST OR REMAINED STABLE FROM BASELINE IN THE 6-MINUTE WALK DISTANCE TEST AND WHO FC HAS IMPROVED OR REMAINED STABLE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRIENTINE

## Products Affected

- *trientine*

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	WILSONS DISEASE: INITIAL: KNOWN FAMILY HISTORY OF WILSONS DISEASE OR PHYSICAL EXAMINATION CONSISTENT WITH WILSONS DISEASE. CONFIRMATION OF ONE OF THE FOLLOWING: 1) PLASMA COPPER-PROTEIN CERULOPLASMIN LESS THAN 20 MG/DL, 2) LIVER BIOPSY POSITIVE FOR AN ABNORMALLY HIGH CONCENTRATION OF COPPER (GREATER THAN 250 MCG/G DRY WEIGHT) OR THE PRESENCE OF KAYSER-FLEISCHER RINGS, OR 3) CONFIRMATION BY GENETIC TESTING FOR ATP7B MUTATIONS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	WILSONS DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST OR GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 12 MONTHS, RENEWAL: LIFETIME.
<b>Other Criteria</b>	WILSONS DISEASE: INITIAL: TRIAL OF OR CONTRAINDICATION TO FORMULARY VERSION OF PENICILLAMINE (DEPEN). RENEWAL: PATIENT CONTINUES TO BENEFIT FROM THE MEDICATION.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRIFLURIDINE/TIPIRACIL

---

## Products Affected

- LONSURF ORAL TABLET 15-6.14 MG, 20-8.19 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# TRIPTORELIN-TRELSTAR

---

## Products Affected

- TRELSTAR INTRAMUSCULAR  
SUSPENSION FOR  
RECONSTITUTION

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS.
<b>Other Criteria</b>	THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TRIPTORELIN-TRIPTODUR

---

## Products Affected

- TRIPTODUR

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	CENTRAL PRECOCIOUS PUBERTY (CPP): INITIAL: FEMALES: ELEVATED LEVELS OF FOLLICLE-STIMULATING HORMONE (FSH) GREATER THAN 4.0 MIU/ML AND LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVELS OF FSH GREATER THAN 5.0 MIU/ML AND LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	CPP: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.
<b>Coverage Duration</b>	INITIAL/RENEWAL: 12 MONTHS.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Other Criteria</b>	<p>CPP: INITIAL: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR BREAST DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) DOCUMENTATION OF PUBERTAL STAGING USING THE TANNER SCALE FOR GENITAL DEVELOPMENT (STAGE 2 OR ABOVE) AND PUBIC HAIR GROWTH (STAGE 2 OR ABOVE). RENEWAL: 1) TANNER SCALE STAGING AT INITIAL DIAGNOSIS OF CPP HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) PATIENT HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# TUCATINIB

---

## Products Affected

- TUKYSA ORAL TABLET 150 MG, 50 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# UBROGEPANT

---

## Products Affected

- UBRELVY

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	ACUTE MIGRAINE TREATMENT: INITIAL: TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN). RENEWAL: 1) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR 2) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# UPADACITINIB

---

## Products Affected

- RINVOQ

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: RA, AS, NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. AD: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST, OR IMMUNOLOGIST. UC: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.

PA Criteria	Criteria Details
<b>Other Criteria</b>	<p>INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD. ATOPIC DERMATITIS: 1) ATOPIC DERMATITIS COVERING AT LEAST 10 PERCENT OF BODY SURFACE AREA OR ATOPIC DERMATITIS AFFECTING THE FACE, HEAD, NECK, HANDS, FEET, GROIN, OR INTERTRIGINOUS AREAS, 2) INTRACTABLE PRURITUS OR CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, 3) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING: TOPICAL CORTICOSTEROID, TOPICAL CALCINEURIN INHIBITOR, TOPICAL PDE4 INHIBITOR, OR TOPICAL JAK INHIBITOR, AND 4) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGIC/JAK INHIBITOR FOR THE TREATMENT OF ATOPIC DERMATITIS. UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPYRINE, METHOTREXATE, OR MESALAMINE. NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID (NON-STEROIDAL ANTI-INFLAMMATORY DRUG). RENEWAL: RA, PSA, AS, NR-AXSPA: CONTINUES TO BENEFIT FROM THE MEDICATION. ATOPIC DERMATITIS: 1) IMPROVEMENT WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGIC/JAK INHIBITOR FOR THE TREATMENT OF ATOPIC DERMATITIS.</p>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# USTEKINUMAB

## Products Affected

- STELARA

PA Criteria	Criteria Details
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, OR FACE.
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
<b>Coverage Duration</b>	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
<b>Other Criteria</b>	INITIAL: PSA: TRIAL OF OR CONTRAINDICATION TO ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG). PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A PUVA (PHOTOTHERAPY ULTRAVIOLET LIGHT A), UVB (ULTRAVIOLET LIGHT B), TOPICAL CORTICOSTEROIDS, CALCIPOTRIENE, ACITRETIN, METHOTREXATE, OR CYCLOSPORINE. CD, UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPYRINE, METHOTREXATE, OR MESALAMINE. RENEWAL: PSA, PSO: CONTINUES TO BENEFIT FROM THE MEDICATION.



<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# USTEKINUMAB IV

---

## Products Affected

- STELARA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.
Coverage Duration	2 MONTHS
Other Criteria	CD, UC: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL THERAPY SUCH AS A CORTICOSTEROID (E.G., BUDESONIDE, METHYLPREDNISOLONE), AZATHIOPRINE, MERCAPTOPYRINE, METHOTREXATE, OR MESALAMINE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# VALBENZINE

---

## Products Affected

- INGREZZA
- INGREZZA INITIATION PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	TARDIVE DYSKINESIA (TD): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR MOVEMENT DISORDER SPECIALIST.
Coverage Duration	12 MONTHS
Other Criteria	TD: 1) PRIOR HISTORY OF USING AGENTS THAT CAUSE TARDIVE DYSKINESIA, AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: AUSTEDO.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# VANDETANIB

---

## Products Affected

- CAPRELSA ORAL TABLET 100 MG, 300 MG

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	CURRENTLY STABLE ON CAPRELSA REQUIRES NO EXTRA CRITERIA.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# VEMURAFENIB

---

## Products Affected

- ZELBORAF

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	MELANOMA: ZELBORAF WILL BE USED ALONE OR IN COMBINATION WITH COTELLIC
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# VENETOCLAX

---

## Products Affected

- VENCLEXTA ORAL TABLET 10 MG, 100 MG, 50 MG
- VENCLEXTA STARTING PACK

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# VESTRONIDASE ALFA VJBK

## Products Affected

- MEPSEVII

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	MUCOPOLYSACCHARIDOSIS VII (MPS VII): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN GENETIC OR METABOLIC DISORDERS.
Coverage Duration	INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.
Other Criteria	MPS VII: INITIAL: 1) PATIENT HAS NOT UNDERGONE SUCCESSFUL BONE MARROW OR STEM CELL TREATMENT FOR MPS VII, 2) LIMITATION IN MOBILITY, BUT REMAINS SUFFICIENTLY AMBULATORY, AND 3) DIAGNOSIS OF MPS VII CONFIRMED BY ALL OF THE FOLLOWING: A) URINARY GAG (GLYCOSAMINOGLYCAN) LEVEL IS GREATER THAN THREE TIMES THE UPPER LEVEL OF NORMAL BASED ON THE LABORATORY ASSAY, B) BETA-GLUCURONIDASE ENZYME ACTIVITY DEFICIENCY OR GENETIC TESTING, AND C) ONE OF THE FOLLOWING CLINICAL SIGNS OF MPS VII: ENLARGED LIVER AND SPLEEN, JOINT LIMITATIONS, AIRWAY OBSTRUCTIONS OR PULMONARY DYSFUNCTION. RENEWAL: IMPROVED, MAINTAINED, OR DEMONSTRATED A LESS THAN EXPECTED DECLINE IN AMBULATORY ABILITY FROM BASELINE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No



# VIGABATRIN

---

## Products Affected

- *vigabatrin*
- *vigadrone*

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	REFRACTORY COMPLEX PARTIAL SEIZURES (CPS), INFANTILE SPASMS: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	CPS: 1) TRIAL OF OR CONTRAINDICATION TO TWO ANTIEPILEPTIC AGENTS AND 2) BENEFITS OUTWEIGH THE POTENTIAL FOR VISION LOSS. INFANTILE SPASMS: BENEFITS OUTWEIGH THE POTENTIAL FOR VISION LOSS.
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# VISMODEGIB

---

## Products Affected

- ERIVEDGE

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

# VORICONAZOLE SUSPENSION

---

## Products Affected

- *voriconazole oral suspension for reconstitution*

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	CANDIDA INFECTIONS: 3 MOS. ALL OTHER INDICATIONS: 6 MOS.
Other Criteria	CANDIDA INFECTIONS: TRIAL OF OR CONTRAINDICATION TO FLUCONAZOLE. ALL INDICATIONS: INABILITY TO SWALLOW TABLETS OR AN INDICATION FOR ESOPHAGEAL CANDIDIASIS. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# ZANUBRUTINIB

---

## Products Affected

- BRUKINSA

<b>PA Criteria</b>	<b>Criteria Details</b>
<b>Exclusion Criteria</b>	
<b>Required Medical Information</b>	
<b>Age Restrictions</b>	
<b>Prescriber Restrictions</b>	
<b>Coverage Duration</b>	12 MONTHS
<b>Other Criteria</b>	
<b>Indications</b>	All FDA-approved Indications.
<b>Off Label Uses</b>	
<b>Part B Prerequisite</b>	No

## INDEX

<i>abiraterone</i> .....	8	BORTEZOMIB INTRAVENOUS	
ACTEMRA.....	387, 389	RECON SOLN.....	51
ACTEMRA ACTPEN.....	389	BOSULIF ORAL TABLET 100 MG,	
ACTHAR.....	73	400 MG, 500 MG.....	54
ACTIMMUNE.....	199	BRAFTOVI ORAL CAPSULE 75 MG	119
ADAKVEO.....	75	BRUKINSA.....	428
ADEMPAS.....	318	<i>butalbital-acetaminop-caff-cod</i> .....	162
AJOVY AUTOINJECTOR.....	143	<i>butalbital-acetaminophen oral tablet 50-</i>	
AJOVY SYRINGE.....	143	<i>325 mg</i> .....	162
ALECENSA.....	16	<i>butalbital-acetaminophen-caff oral</i>	
ALTRENO.....	394	<i>capsule 50-325-40 mg</i> .....	162
ALUNBRIG ORAL TABLET 180 MG,		<i>butalbital-acetaminophen-caff oral tablet</i>	162
30 MG, 90 MG.....	55	<i>butalbital-aspirin-caffeine</i> .....	162
ALUNBRIG ORAL TABLETS,DOSE		CABLIVI INJECTION KIT.....	62
PACK.....	55	CABOMETYX ORAL TABLET 20	
<i>alyq</i> .....	279	MG, 40 MG, 60 MG.....	59
<i>amabelz</i> .....	165	CALQUENCE.....	10
<i>ambrisentan</i> .....	18	CALQUENCE (ACALABRUTINIB	
<i>apomorphine</i> .....	22	MAL).....	10
<i>armodafinil</i> .....	249	CAPRELSA ORAL TABLET 100 MG,	
<i>ascomp with codeine</i> .....	162	300 MG.....	420
AUBAGIO.....	376	<i>carbinoxamine maleate oral liquid</i> .....	170
AUSTEDO ORAL TABLET 12 MG, 6		<i>carbinoxamine maleate oral tablet 4 mg..</i>	170
MG, 9 MG.....	94	<i>carglumic acid</i> .....	64
AVONEX INTRAMUSCULAR PEN		CAYSTON.....	36
INJECTOR KIT.....	196	CERDELGA.....	113
AVONEX INTRAMUSCULAR		<i>chlorzoxazone oral tablet 250 mg, 500 mg</i>	
SYRINGE KIT.....	196	.....	169
AVSOLA.....	191	CIMZIA.....	66
AYVAKIT.....	32	CIMZIA POWDER FOR RECONST....	66
BALVERSA ORAL TABLET 3 MG, 4		CINQAIR.....	311
MG, 5 MG.....	125	CINRYZE.....	56
<i>bendamustine intravenous recon soln</i> .....	43	<i>clemastine oral tablet 2.68 mg</i> .....	158
BENDEKA.....	43	<i>codeine-butalbital-asa-caff</i> .....	162
BENLYSTA INTRAVENOUS.....	40	<i>colchicine oral tablet</i> .....	72
BENLYSTA SUBCUTANEOUS.....	40	COMETRIQ ORAL CAPSULE 100	
BESREMI.....	331	MG/DAY(80 MG X1-20 MG X1), 140	
<i>betaine</i> .....	46	MG/DAY(80 MG X1-20 MG X3), 60	
BETASERON SUBCUTANEOUS KIT		MG/DAY (20 MG X 3/DAY).....	58
.....	197	COPAXONE SUBCUTANEOUS	
<i>bexarotene</i> .....	49	SYRINGE 20 MG/ML, 40 MG/ML.....	151
<i>bortezomib injection recon soln 1 mg, 2.5</i>		COPIKTRA.....	105
<i>mg</i> .....	51	CORTROPHIN GEL.....	73

COSENTYX (2 SYRINGES).....	340	ELIGARD (4 MONTH).....	218
COSENTYX PEN (2 PENS).....	340	ELIGARD (6 MONTH).....	218
COSENTYX SUBCUTANEOUS		EMGALITY PEN.....	145
SYRINGE 75 MG/0.5 ML.....	340	EMGALITY SYRINGE	
COTELLIC.....	71	SUBCUTANEOUS SYRINGE 120	
<i>cyclobenzaprine oral tablet 10 mg, 5 mg..</i>	169	MG/ML, 300 MG/3 ML (100 MG/ML	
<i>cyproheptadine.....</i>	170	X 3).....	145
CYRAMZA.....	308	ENBREL.....	128
CYSTADROPS.....	77	ENBREL MINI.....	128
CYSTARAN.....	77	ENBREL SURECLICK.....	128
<i>dalfampridine.....</i>	80	ENDARI.....	224
DANYELZA.....	251	ENSPRYNG.....	337
DARZALEX.....	81	ENTADFI.....	138
DARZALEX FASPRO.....	82	EPCLUSA ORAL PELLETS IN	
DAURISMO ORAL TABLET 100		PACKET 150-37.5 MG, 200-50 MG.....	351
MG, 25 MG.....	150	EPCLUSA ORAL TABLET.....	351
<i>deferasirox.....</i>	87	EPIDIOLEX.....	61
<i>deferiprone.....</i>	89	<i>epoprostenol (glycine).....</i>	124
<i>deferoxamine.....</i>	91	ERBITUX.....	68
DIACOMIT ORAL CAPSULE 250		ERIVEDGE.....	426
MG, 500 MG.....	362	ERLEADA.....	21
DIACOMIT ORAL POWDER IN		<i>erlotinib oral tablet 100 mg, 150 mg, 25</i>	
PACKET 250 MG, 500 MG.....	362	<i>mg.....</i>	126
<i>diclofenac sodium topical gel 3 %.....</i>	95	ESBRIET ORAL CAPSULE.....	298
<i>diclofenac sodium topical solution in</i>		<i>estradiol oral.....</i>	165
<i>metered-dose pump.....</i>	96	<i>estradiol transdermal patch semiweekly...165</i>	
<i>dimethyl fumarate oral capsule, delayed</i>		<i>estradiol transdermal patch weekly.....165</i>	
<i>release(drlec) 120 mg, 120 mg (14)- 240</i>		<i>estradiol-norethindrone acet oral tablet</i>	
<i>mg (46), 240 mg.....</i>	97	<i>0.5-0.1 mg.....</i>	165
<i>diphenhydramine hcl oral elixir.....</i>	171	EVENITY.....	329
<i>diphenoxylate-atropine.....</i>	173	<i>everolimus (antineoplastic) oral tablet 10</i>	
<i>dipyridamole oral.....</i>	163	<i>mg, 2.5 mg, 5 mg, 7.5 mg.....</i>	131
<i>disopyramide phosphate oral capsule.....</i>	164	<i>everolimus (antineoplastic) oral tablet for</i>	
DOPTELET (10 TAB PACK).....	33	<i>suspension.....</i>	131
DOPTELET (15 TAB PACK).....	33	EVRYSDI.....	323
DOPTELET (30 TAB PACK).....	33	EXKIVITY.....	245
<i>dotti.....</i>	165	EXONDYS-51.....	130
<i>dronabinol.....</i>	100	FABRAZYME.....	15
<i>droxidopa.....</i>	101	FARYDAK.....	275
DUAVEE.....	165	FASENRA.....	44
DUPIXENT PEN.....	102	FASENRA PEN.....	44
DUPIXENT SYRINGE.....	102	<i>fentanyl citrate buccal lozenge on a</i>	
EGRIFTA SV.....	379	<i>handle.....</i>	134
ELIGARD.....	218	FERRIPROX (2 TIMES A DAY).....	89
ELIGARD (3 MONTH).....	218	FERRIPROX ORAL SOLUTION.....	89

<i>fingolimod</i> .....	140	HUMIRA(CF) PEN CROHNS-UC-HS .	12
FINTEPLA.....	133	HUMIRA(CF) PEN PEDIATRIC UC...	12
FORTEO.....	377	HUMIRA(CF) PEN PSOR-UV-ADOL	
FOTIVDA.....	386	HS.....	12
FULPHILA.....	286	IBRANCE.....	273
FYARRO.....	348	<i>ibuprofen-famotidine</i> .....	182
<i>fyavolv</i> .....	165	<i>icatibant</i> .....	183
FYLNETRA.....	289	ICLUSIG.....	301
GALAFOLD.....	242	IDHIFA.....	118
GAMIFANT.....	116	ILARIS (PF).....	60
GATTEX 30-VIAL.....	372	ILUMYA.....	384
GAVRETO.....	304	<i>imatinib oral tablet 100 mg, 400 mg</i> .....	185
GILENYA ORAL CAPSULE 0.25 MG		IMBRUVICA ORAL CAPSULE 140	
.....	140	MG, 70 MG.....	181
GILOTRIF.....	14	IMBRUVICA ORAL SUSPENSION...	181
GIVLAARI.....	149	IMBRUVICA ORAL TABLET.....	181
<i>glatiramer subcutaneous syringe 20</i>		IMJUDO.....	402
<i>mg/ml, 40 mg/ml</i> .....	151	IMLYGIC INJECTION SUSPENSION	
<i>glatopa subcutaneous syringe 20 mg/ml,</i>		10EXP6 (1 MILLION) PFU/ML.....	366
<i>40 mg/ml</i> .....	151	IMPAVIDO.....	244
<i>glyburide</i> .....	167	INBRIJA INHALATION CAPSULE,	
<i>glyburide micronized</i> .....	167	W/INHALATION DEVICE.....	223
<i>glyburide-metformin</i> .....	167	<i>indomethacin oral capsule 25 mg, 50 mg</i> ..	174
GRANIX.....	369	<i>indomethacin oral capsule, extended</i>	
HAEGARDA SUBCUTANEOUS		<i>release</i> .....	174
RECON SOLN 2,000 UNIT, 3,000		INFLECTRA.....	193
UNIT.....	57	<i>infliximab</i> .....	187
HARVONI ORAL PELLETS IN		INGREZZA.....	419
PACKET 33.75-150 MG, 45-200 MG...	211	INGREZZA INITIATION PACK.....	419
HARVONI ORAL TABLET.....	211	INLYTA ORAL TABLET 1 MG, 5	
HERCEPTIN HYLECTA.....	396	MG.....	34
HERZUMA.....	400	INQOVI.....	86
HETLIOZ.....	367	INREBIC.....	132
HETLIOZ LQ.....	367	INTRON A INJECTION RECON	
HUMIRA PEN.....	12	SOLN.....	195
HUMIRA PEN CROHNS-UC-HS		IRESSA.....	147
START.....	12	<i>itraconazole oral solution</i> .....	201
HUMIRA PEN PSOR-UEVITS-ADOL		JAKAFI.....	333
HS.....	12	<i>javygtor oral tablet, soluble</i> .....	335
HUMIRA SUBCUTANEOUS		JEMPERLI.....	99
SYRINGE KIT 40 MG/0.8 ML.....	12	<i>jinteli</i> .....	165
HUMIRA(CF).....	12	JUXTAPID ORAL CAPSULE 10 MG,	
HUMIRA(CF) PEDI CROHNS		20 MG, 30 MG, 40 MG, 5 MG, 60 MG	227
STARTER.....	12	JYNARQUE ORAL TABLET.....	393
HUMIRA(CF) PEN.....	12		

JYNARQUE ORAL TABLETS, SEQUENTIAL.....	393	<i>lidocaine topical ointment</i> .....	225
KALYDECO.....	202	<i>lidocaine-prilocaine topical cream</i> .....	226
KANJINTI.....	397	LONSURF ORAL TABLET 15-6.14 MG, 20-8.19 MG.....	408
KANUMA.....	338	LORBRENA ORAL TABLET 100 MG, 25 MG.....	230
KERENDIA.....	139	LUMAKRAS.....	361
KESIMPTA PEN.....	262	LUNSUMIO.....	246
<i>ketorolac injection cartridge 15 mg/ml</i> .....	168	LUPRON DEPOT.....	219
<i>ketorolac injection solution 15 mg/ml, 30 mg/ml, 30 mg/ml (1 ml)</i> .....	168	LUPRON DEPOT (3 MONTH).....	219
<i>ketorolac injection syringe 15 mg/ml, 30 mg/ml</i> .....	168	LUPRON DEPOT (4 MONTH).....	219
<i>ketorolac intramuscular solution</i> .....	168	LUPRON DEPOT (6 MONTH).....	219
<i>ketorolac intramuscular syringe</i> .....	168	LUPRON DEPOT-PED (3 MONTH) INTRAMUSCULAR SYRINGE KIT 30 MG.....	221
<i>ketorolac oral</i> .....	168	LUPRON DEPOT-PED INTRAMUSCULAR KIT 11.25 MG, 15 MG.....	221
KEVZARA.....	336	LYBALVI.....	263
KEYTRUDA.....	292	<i>lyllana</i> .....	165
KIMMTRAK.....	370	LYNPARZA.....	264
KINERET.....	20	LYTGOBI.....	144
KISQALI FEMARA CO-PACK ORAL TABLET 200 MG/DAY(200 MG X 1)- 2.5 MG, 400 MG/DAY(200 MG X 2)- 2.5 MG, 600 MG/DAY(200 MG X 3)- 2.5 MG.....	314	MARGENZA.....	234
KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3).....	314	MAVENCLAD (10 TABLET PACK)....	69
KORLYM.....	241	MAVENCLAD (4 TABLET PACK).....	69
KOSELUGO ORAL CAPSULE 10 MG, 25 MG.....	346	MAVENCLAD (5 TABLET PACK).....	69
KRAZATI.....	11	MAVENCLAD (6 TABLET PACK).....	69
KYNMOBI SUBLINGUAL FILM 10 MG, 10-15-20-25-30 MG, 15 MG, 20 MG, 25 MG, 30 MG.....	23	MAVENCLAD (7 TABLET PACK).....	69
<i>lanreotide</i> .....	208	MAVENCLAD (8 TABLET PACK).....	69
<i>lapatinib</i> .....	209	MAVENCLAD (9 TABLET PACK).....	69
<i>lenalidomide</i> .....	212	MAVYRET ORAL TABLET.....	152
LENVIMA.....	213	MAYZENT ORAL TABLET 0.25 MG, 1 MG, 2 MG.....	347
<i>leuprolide (3 month)</i> .....	217	MAYZENT STARTER(FOR 1MG MAINT).....	347
<i>leuprolide subcutaneous kit</i> .....	215	MAYZENT STARTER(FOR 2MG MAINT).....	347
<i>lidocaine hcl mucous membrane solution 4 % (40 mg/ml)</i> .....	225	<i>megestrol oral suspension 400 mg/10 ml (40 mg/ml)</i> .....	175
<i>lidocaine topical adhesive patch,medicated 5%</i> .....	225	<i>megestrol oral tablet</i> .....	175
		MEKINIST ORAL TABLET 0.5 MG, 2 MG.....	395
		MEKTOVI.....	50
		MEPSEVII.....	423



<i>methocarbamol oral tablet 500 mg, 750 mg</i> .....	169	ORENCIA (WITH MALTOSE).....	3
<i>miglustat</i> .....	243	ORENCIA CLICKJECT.....	5
<i>mimvey</i> .....	165	ORFADIN ORAL CAPSULE 20 MG.....	257
<i>modafinil oral tablet 100 mg, 200 mg</i> .....	249	ORFADIN ORAL SUSPENSION.....	257
<i>morphine concentrate oral solution</i> .....	157	ORGOVYX.....	310
MVASI.....	47	ORLISSA ORAL TABLET 150 MG, 200 MG.....	109
NATPARA.....	276	ORKAMBI ORAL GRANULES IN PACKET.....	231
NERLYNX.....	252	ORKAMBI ORAL TABLET.....	231
NEULASTA.....	283	OTEZLA.....	24
NEULASTA ONPRO.....	288	OTEZLA STARTER.....	24
NINLARO.....	204	OXLUMO.....	232
<i>nitisinone</i> .....	257	<i>oxycodone oral concentrate</i> .....	157
NIVESTYM.....	135	PALYNZIQ.....	290
NORDITROPIN FLEXPRO.....	355	<i>paroxetine hcl</i> .....	176
<i>norethindrone ac-eth estradiol oral tablet 0.5-2.5 mg-mcg, 1-5 mg-mcg</i> .....	165	PEMAZYRE.....	293
NOXAFIL ORAL SUSP,DELAYED RELEASE FOR RECON.....	303	<i>penicillamine oral tablet</i> .....	294
NOXAFIL ORAL SUSPENSION.....	302	<i>phenobarbital</i> .....	177
NPLATE.....	328	PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2).....	17
NUBEQA.....	83	<i>pirfenidone oral capsule</i> .....	298
NUCALA SUBCUTANEOUS AUTO- INJECTOR.....	235	<i>pirfenidone oral tablet 267 mg, 534 mg, 801 mg</i> .....	298
NUCALA SUBCUTANEOUS RECON SOLN.....	235	PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML.....	198
NUCALA SUBCUTANEOUS SYRINGE 100 MG/ML, 40 MG/0.4 ML.....	235	PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML, 63 MCG/0.5 ML- 94 MCG/0.5 ML.....	198
NUPLAZID.....	297	POMALYST.....	300
NURTEC ODT.....	316	<i>posaconazole</i> .....	302
NYVEPRIA.....	284	PREMARIN ORAL.....	165
OALIVA.....	260	PREMPHASE.....	165
OCREVUS.....	261	PREMPRO.....	165
ODOMZO.....	359	PREVYMIS INTRAVENOUS SOLUTION 240 MG/12 ML, 480 MG/24 ML.....	214
OFEV.....	254	PREVYMIS ORAL.....	214
OGIVRI.....	398	PROMACTA ORAL POWDER IN PACKET 12.5 MG, 25 MG.....	115
OLUMIANT.....	37	PROMACTA ORAL TABLET 12.5 MG, 25 MG, 50 MG, 75 MG.....	115
ONGENTYS.....	270		
ONTRUZANT.....	399		
ONUREG.....	35		
OPDIVO.....	258		
OPDUALAG.....	259		
OPSUMIT.....	233		
ORENCIA.....	5		

<i>promethazine injection solution</i> .....	159	SIGNIFOR.....	277
<i>promethazine oral</i> .....	159	SIKLOS ORAL TABLET 100 MG.....	180
<i>promethazine rectal</i> .....	159	<i>sildenafil (pulm.hypertension)</i>	
<i>promethegan</i> .....	159	<i>intravenous</i> .....	281
<i>pyrimethamine</i> .....	306	<i>sildenafil (pulm.hypertension) oral tablet</i>	279
QINLOCK.....	320	SIRTURO.....	39
<i>quinine sulfate</i> .....	307	SKYRIZI.....	321
QULIPTA.....	31	<i>sodium oxybate</i> .....	349
RADICAVA.....	106	SOMATULINE DEPOT	
RAVICTI.....	154	SUBCUTANEOUS SYRINGE 120	
REGRANEX.....	38	MG/0.5 ML, 60 MG/0.2 ML, 90 MG/0.3	
RELEUKO.....	136	ML.....	208
RELISTOR ORAL.....	239	SOMAVERT.....	291
RELISTOR SUBCUTANEOUS		<i>sorafenib</i> .....	360
SOLUTION.....	238	SPRAVATO.....	127
RELISTOR SUBCUTANEOUS		SPRYCEL ORAL TABLET 100 MG,	
SYRINGE 12 MG/0.6 ML, 8 MG/0.4		140 MG, 20 MG, 50 MG, 70 MG, 80	
ML.....	238	MG.....	85
RENFLEXIS.....	189	STELARA.....	416, 418
RETACRIT INJECTION SOLUTION		STIVARGA.....	309
10,000 UNIT/ML, 2,000 UNIT/ML,		STRENSIQ.....	27
20,000 UNIT/2 ML, 20,000 UNIT/ML,		<i>sunitinib</i> .....	363
3,000 UNIT/ML, 4,000 UNIT/ML,		SUNOSI.....	354
40,000 UNIT/ML.....	122	SUPPRELIN LA.....	178
RETEVMO ORAL CAPSULE 40 MG,		SYMDEKO.....	382
80 MG.....	345	SYMLINPEN 120.....	305
REVCOVI.....	111	SYMLINPEN 60.....	305
REZLIDHIA.....	265	SYMPAZAN.....	70
REZUROCK.....	41	SYNAGIS.....	274
RIABNI.....	326	SYNAREL.....	247
RINVOQ.....	414	SYNRIBO.....	266
RITUXAN HYCELA.....	324	TABRECTA.....	63
ROLVEDON.....	108	<i>tadalafil (pulm. hypertension)</i> .....	279
ROZLYTREK ORAL CAPSULE 100		<i>tadalafil oral tablet 2.5 mg, 5 mg</i> .....	364
MG, 200 MG.....	120	TAFINLAR.....	78
RUBRACA.....	332	TAGRISSE.....	271
RUXIENCE.....	327	TAKHZYRO.....	207
RYBREVANT.....	19	TALTZ AUTOINJECTOR.....	205
RYDAPT.....	240	TALTZ SYRINGE.....	205
<i>sajazir</i> .....	183	TALZENNA.....	365
<i>sapropterin oral tablet, soluble</i> .....	335	TASCENSO ODT.....	141
SCSEMBLIX.....	26	TASIGNA ORAL CAPSULE 150 MG,	
<i>scopolamine base</i> .....	161	200 MG, 50 MG.....	253
SEROSTIM SUBCUTANEOUS		<i>tasimelteon</i> .....	367
RECON SOLN 4 MG, 5 MG, 6 MG....	357	TAVALISSE.....	142

TAZVERIK.....	368	UPTRAVI ORAL TABLET 1,000	
TECENTRIQ.....	30	MCG, 1,200 MCG, 1,400 MCG, 1,600	
TECVAYLI.....	371	MCG, 200 MCG, 400 MCG, 600 MCG,	
<i>tencon</i> .....	162	800 MCG.....	342
TEPEZZA.....	375	UPTRAVI ORAL TABLETS,DOSE	
TEPMETKO.....	374	PACK.....	342
<i>testosterone cypionate</i> .....	380	VELCADE.....	51
<i>testosterone enanthate</i> .....	380	VENCLEXTA ORAL TABLET 10	
<i>testosterone transdermal gel in metered-</i>		MG, 100 MG, 50 MG.....	422
<i>dose pump 12.5 mg/ 1.25 gram (1%),</i>		VENCLEXTA STARTING PACK.....	422
<i>20.25 mg/1.25 gram (1.62%)</i> .....	380	VERZENIO.....	7
<i>testosterone transdermal gel in packet 1%</i>		<i>vigabatrin</i> .....	425
<i>(25 mg/2.5gram), 1% (50 mg/5 gram)</i> ...	380	<i>vigadrone</i> .....	425
<i>testosterone transdermal solution in</i>		VIMIZIM.....	114
<i>metered pump w/lapp</i> .....	380	VITRAKVI ORAL CAPSULE 100	
<i>tetrabenazine</i> .....	381	MG, 25 MG.....	210
THALOMID.....	383	VITRAKVI ORAL SOLUTION.....	210
TIBSOVO.....	203	VIZIMPRO.....	79
TIVDAK.....	385	VONJO.....	272
TRACLEER ORAL TABLET.....	52	<i>voriconazole oral suspension for</i>	
TRACLEER ORAL TABLET FOR		<i>reconstitution</i> .....	427
SUSPENSION.....	52	VOSEVI.....	352
TRAZIMERA.....	401	VOTRIENT.....	278
TRELSTAR INTRAMUSCULAR		VUMERITY.....	98
SUSPENSION FOR		WELIREG.....	42
RECONSTITUTION.....	409	XADAGO.....	334
TREMFYA.....	156	XALKORI.....	76
<i>treprostinil sodium</i> .....	405	XELJANZ.....	391
<i>tretinoin</i> .....	394	XELJANZ XR.....	391
<i>trientine</i> .....	407	XERMELO.....	373
TRIKAFTA.....	112	XGEVA.....	93
TRIPTODUR.....	410	XIFAXAN ORAL TABLET 200 MG,	
TRUSELTIQ.....	186	550 MG.....	315
TRUXIMA.....	325	XOLAIR.....	267
TUKYSA ORAL TABLET 150 MG, 50		XOSPATA.....	148
MG.....	412	XPOVIO ORAL TABLET 100	
TURALIO ORAL CAPSULE 200 MG	296	MG/WEEK (50 MG X 2), 40	
TYMLOS.....	1	MG/WEEK (40 MG X 1), 40MG	
TYSABRI.....	250	TWICE WEEK (40 MG X 2), 60	
TYVASO.....	403	MG/WEEK (60 MG X 1), 60MG	
UBRELVY.....	413	TWICE WEEK (120 MG/WEEK), 80	
UDENYCA.....	285	MG/WEEK (40 MG X 2), 80MG	
UPTRAVI INTRAVENOUS.....	342	TWICE WEEK (160 MG/WEEK).....	344
		XTANDI ORAL CAPSULE.....	121

XTANDI ORAL TABLET 40 MG, 80	
MG .....	121
XYOSTED .....	380
XYREM .....	349
YERVOY .....	200
YONSA .....	9
ZARXIO .....	137
<i>zebutal</i> .....	162
ZEJULA .....	256
ZELBORAF .....	421
ZIEXTENZO .....	287
ZIRABEV .....	48
ZOLADEX .....	155
ZTALMY .....	146
ZTLIDO .....	225
ZYDELIG .....	184
ZYKADIA .....	65
ZYNLONTA .....	229